arXiv:1001.3514v2 [math.DG] 28 Nov 2011

On the indices of minimal orbits of Hermann actions

Naoyuki Koike

Abstract

We give a formula to estimate the indices of special (non-totally geodesic) minimal orbits of Hermann actions. Also, we give examples of such minimal orbits of Hermann actions and estimate their indices by using the formula.

1 Introduction

Y. Ohnita [O] gave a formula to calculate the indices (and nullities) of totally geodesic submanifolds in a symmetric space N of compact type and showed that the indices of all Helgason spheres in every simply connected irreducible compact symmetric space are equal to zero, that is, they are stable. By using this index formula, M. S. Tanaka [Ta] determined the stability of all polars and meridians in every simply connected irreducible compact symmetric space. Note that polars and meridians are totally geodesic. By using this index formula, T. Kimura [Ki] determined the stability of all totally geodesic singular orbits of all cohomogeneity one actions on every simply connected irreducible compact symmetric space. Also, by using this index formula, T. Kimura and M. S. Tanaka [MK] determined the stability of all maximal totally geodesic submanifolds in every simply connected irreducible compact symmetric space of rank two. Let $N = G/K$ be a symmetrc space of compact type and H be a symmetric subgroup of G (i.e., $(\mathrm{Fix}\,\tau)_0 \subset H \subset \mathrm{Fix}\,\tau$ for some involution τ of G), where $\mathrm{Fix}\,\tau$ is the fixed point group of τ and $(\mathrm{Fix}\,\tau)_0$ is the identity component of $\mathrm{Fix}\,\tau$. The natural action of H on N is called a *Hermann action* (see [HPTT], [Kol]). Let θ be an involution of G with $(\mathrm{Fix}\,\theta)_0 \subset K \subset \mathrm{Fix}\,\theta$. According to [Co], we may assume that $\theta \circ \tau = \tau \circ \theta$ by replacing H to a suitable conjugate group of H if necessary except for the following three Hermann action:

(i) $Sp(p+q) \curvearrowright SU(2p+2q)/S(U(2p-1) \times U(2q+1)) \quad (p \geq q+2)$,
(ii) $U(p+q+1) \curvearrowright Spin(2p+2q+2)/Spin(2p+1) \times_{\mathbf{Z}_2} Spin(2q+1) \quad (p \geq q+1)$,
(iii) $Spin(3) \times_{\mathbf{Z}_2} \times Spin(5) \curvearrowright Spin(8)/\omega(Spin(3) \times_{\mathbf{Z}_2} Spin(5))$,

where ω is the triality automorphism of $Spin(8)$. Here we note that we remove transitive Hermann actions.

Assumption. In the sequel, we assume that $\theta \circ \tau = \tau \circ \theta$.

Let $\mathfrak{g}, \mathfrak{k}$ and $\mathfrak{h}$ be the Lie algebras of G, K and H, respectively. Denote the involutions of $\mathfrak{g}$ induced form θ and τ by the same symbols θ and τ, respectively. Set $\mathfrak{p} := \mathrm{Ker}(\theta + \mathrm{id})$ and $\mathfrak{q} := \mathrm{Ker}(\tau + \mathrm{id})$. The vector space $\mathfrak{p}$ is identified with $T_{eK}(G/K)$, where e is the identity element of G. Take a maximal abelian subspace $\mathfrak{b}$ of $\mathfrak{p} \cap \mathfrak{q}$. For each $\beta \in \mathfrak{b}^*$, we set $\mathfrak{p}_\beta := \{X \in \mathfrak{p} \,|\, \mathrm{ad}(b)^2(X) = -\beta(b)^2 X \ (\forall\, b \in \mathfrak{b})\}$ and $\triangle' := \{\beta \in \mathfrak{b}^* \setminus \{0\} \,|\, \mathfrak{p}_\beta \neq \{0\}\}$. This set $\triangle'$ is a root system. Let $\Pi' = \{\beta_1, \cdots, \beta_r\}$ be the simple root system of the positive root system $\triangle'_+$ of $\triangle'$ under a lexicographic ordering of $\mathfrak{b}^*$ and δ be the highest root of $\triangle'_+$. Set ${\triangle'}^V_+ := \{\beta \in \triangle'_+ \,|\, \mathfrak{p}_\beta \cap \mathfrak{q} \neq \{0\}\}$ and ${\triangle'}^H_+ := \{\beta \in \triangle'_+ \,|\, \mathfrak{p}_\beta \cap \mathfrak{h} \neq \{0\}\}$. Define a subset $\widetilde{C}$ of $\mathfrak{b}$ by

$$\widetilde{C} := \{b \in \mathfrak{b} \,|\, 0 < \beta(b) < \pi \, (\forall\, \beta \in {\triangle'}^V_+), \ -\frac{\pi}{2} < \beta(b) < \frac{\pi}{2} \, (\forall\, \beta \in {\triangle'}^H_+)\}.$$

Set $C := \mathrm{Exp}(\widetilde{C})$, where Exp is the exponential map of G/K at eK. Each principal H-orbit passes through only one point of C and each singular H-orbit passes through only one point of $\mathrm{Exp}(\partial\widetilde{C})$. For each simplex σ of $\partial\widetilde{C}$, only one minimal singular H-orbit through $\mathrm{Exp}(\sigma)$ exists. See proofs of Theorems A and B in [Koi2] (also [I]) about this fact. Denote by $D(H)$ the set of all equivalence classes of finite dimensional irreducible complex representations of H and $\rho_{G/H}$ the isotropy representation of G/H, that is, $\rho_{G/H}(h) := \mathrm{Ad}_G(h)|_{\mathfrak{q}}$ $(h \in H)$. Denote by the same symbol $\rho_{G/H}$ the complexification of $\rho_{G/H}$ and μ the equivalence class of the complexification. Denote by $B_{\mathfrak{g}}$ the Killing form of $\mathfrak{g}$. For $\beta \in \triangle'_+$, we set $\beta = \sum_{i=1}^{r} n_i^\beta \beta_i$, $m_\beta := \dim \mathfrak{p}_\beta$, $m_\beta^V := \dim(\mathfrak{p}_\beta \cap \mathfrak{q})$ and $m_\beta^H := \dim(\mathfrak{p}_\beta \cap \mathfrak{h})$. For $Z_0 \in \widetilde{C}$, we consider the following two kinds of conditions on Z_0 as follows:

$$\text{(I)} \qquad \begin{aligned} &\{\beta(Z_0) \,|\, \beta \in {\triangle'}^V_+ \cap {\triangle'}^H_+\} \subset \{0, \frac{\pi}{2}\} \\ &\& \ \{\beta(Z_0) \,|\, \beta \in {\triangle'}^V_+ \setminus {\triangle'}^H_+\} \subset \{0, \frac{\pi}{3}, \frac{2}{3}\pi, \pi\} \\ &\& \ \{\beta(Z_0) \,|\, \beta \in {\triangle'}^H_+ \setminus {\triangle'}^V_+\} \subset \{\pm\frac{\pi}{2}, \pm\frac{\pi}{6}\} \ \& \\ &\sum_{\beta \in {\triangle'}^V_+ \setminus {\triangle'}^H_+ \text{ s.t. } \beta(Z_0) = \frac{\pi}{3}} n_i^\beta m_\beta + \sum_{\beta \in {\triangle'}^H_+ \setminus {\triangle'}^V_+ \text{ s.t. } \beta(Z_0) = -\frac{\pi}{6}} n_i^\beta m_\beta \\ &= \sum_{\beta \in {\triangle'}^V_+ \setminus {\triangle'}^H_+ \text{ s.t. } \beta(Z_0) = \frac{2}{3}\pi} n_i^\beta m_\beta + \sum_{\beta \in {\triangle'}^H_+ \setminus {\triangle'}^V_+ \text{ s.t. } \beta(Z_0) = \frac{\pi}{6}} n_i^\beta m_\beta. \end{aligned}$$

and

$$
\text{(II)}\quad
\begin{aligned}
&\{\beta(Z_0)\,|\,\beta\in\triangle'^V_+\cap\triangle'^H_+\}\subset\{0,\frac{\pi}{4},\frac{\pi}{2}\}\\
&\&\ \{\beta(Z_0)\,|\,\beta\in\triangle'^V_+\setminus\triangle'^H_+\}\subset\{0,\frac{\pi}{4},\frac{3}{4}\pi,\pi\}\\
&\&\ \{\beta(Z_0)\,|\,\beta\in\triangle'^H_+\setminus\triangle'^V_+\}\subset\{\pm\frac{\pi}{2},\pm\frac{\pi}{4}\}\ \&\\
&\sum_{\beta\in\triangle'^V_+\cap\triangle'^H_+\text{ s.t. }\beta(Z_0)=\frac{\pi}{4}} n_i^\beta m_\beta^V+\sum_{\beta\in\triangle'^V_+\setminus\triangle'^H_+\text{ s.t. }\beta(Z_0)=\frac{\pi}{4}} n_i^\beta m_\beta\\
&\qquad\qquad+\sum_{\beta\in\triangle'^H_+\setminus\triangle'^V_+\text{ s.t. }\beta(Z_0)=-\frac{\pi}{4}} n_i^\beta m_\beta\\
&=\sum_{\beta\in\triangle'^V_+\cap\triangle'^H_+\text{ s.t. }\beta(Z_0)=\frac{\pi}{4}} n_i^\beta m_\beta^H+\sum_{\beta\in\triangle'^V_+\setminus\triangle'^H_+\text{ s.t. }\beta(Z_0)=\frac{3}{4}\pi} n_i^\beta m_\beta\\
&\qquad\qquad+\sum_{\beta\in\triangle'^H_+\setminus\triangle'^V_+\text{ s.t. }\beta(Z_0)=\frac{\pi}{4}} n_i^\beta m_\beta.
\end{aligned}
$$

Set $\triangle'^V_{Z_0}:=\{\beta\in\triangle'^V_+\,|\,\beta(Z_0)=0\text{ or }\pi\}$ and $\triangle'^H_{Z_0}:=\{\beta\in\triangle'^H_+\,|\,\beta(Z_0)=\pm\frac{\pi}{2}\}$. Denote by H_{Z_0} the isotropy group of H at $\mathrm{Exp}\,Z_0$. For simplicity, we set $L:=H_{Z_0}$. Denote by $\mathfrak{h}$ (resp. $\mathfrak{h}_{Z_0}$) the Lie algebra of H (resp. L) and $B_{\mathfrak{h}}$ the Killing form of $\mathfrak{h}$. We have a natural reductive decomposition $\mathfrak{h}=\mathfrak{l}+\mathfrak{m}$ (see Step I in the proof of Theorem A). For $Z_0\in\widetilde{C}$ satisfying (I) or (II), $M:=H(\mathrm{Exp}\,Z_0)(=H/L)$ is minimal in G/K and the induced metric on the submanifold M in G/K coincides with the H-invariant metric arising from the restriction $cB_{\mathfrak{h}}|_{\mathfrak{m}\times\mathfrak{m}}$ of some constant-multiple $cB_{\mathfrak{h}}$ of $B_{\mathfrak{h}}$ to $\mathfrak{m}\times\mathfrak{m}$ (see Step II in the proof of Theorem A and the proofs of Propositions in Section 4). Denote by $\rho^S_{Z_0}$ the slice representation of the orbit M at $\mathrm{Exp}\,Z_0$. Under the identification of the normal space $T^\perp_{\mathrm{Exp}\,Z_0}(M)$ of M at $\mathrm{Exp}\,Z_0$ and $\mathfrak{m}^\perp:=(\exp\,Z_0)_*^{-1}(T^\perp_{\mathrm{Exp}\,Z_0}(M))(\subset\mathfrak{p}\cap\mathfrak{q})$, $\rho^S_{Z_0}$ is identified with $(\rho_{G/H}|_L)|_{\mathfrak{m}^\perp}$, where $(\rho_{G/H}|_L)|_{\mathfrak{m}^\perp}$ means the representaion of L with representation space $\mathfrak{m}^\perp$ defined by $((\rho_{G/H}|_L)|_{\mathfrak{m}^\perp})(l))(w):=(\rho_{G/H}(l))(w)$ $(l\in L,\,w\in\mathfrak{m}^\perp)$ in terms of the isotropy representation $\rho_{G/H}$ of G/H. We regard $(\mathfrak{m}^\perp)^{\mathbf{c}}$ as a L-module under the complexification of $(\rho_{G/H}|_L)|_{\mathfrak{m}^\perp}$.

In this paper, we prove the following result.

Theorem A. *Let G/K be an irreducible simply connected symmetric space of compact type, $H\curvearrowright G/K$ be a Hermann action whose codimension is equal to the rank of G/K and Z_0 be an element of the closure of $\widetilde{C}$ satisfying the above condition (I) or (II). Then, the orbit $M:=H(\mathrm{Exp}\,Z_0)$ is a (non-totally geodesic) unstable minimal submanifold and the index $i(M)$ of M is estimated as follows:*

$$\sum_{\lambda\in D_1(H)} m_\lambda\cdot\mathrm{dimHom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}})\le i(M)\le\sum_{\lambda\in D_2(H)} m_\lambda\cdot\dim\mathrm{Hom}_L(V_\lambda,(\mathfrak{m}^\perp)^{\mathbf{c}}).$$

Here $D_1(H) := \{\lambda \in D(H) \,|\, a_\lambda \geq a_\mu\}$ *and* $D_2(H) := \{\lambda \in D(H) \,|\, a_\lambda > a_\mu - || \sum\limits_{\beta \in (\triangle'^V_+ \setminus \triangle'^V_{Z_0}) \cup (\triangle'^H_+ \setminus \triangle'^H_{Z_0})} m_\beta \hat{c}(\beta \otimes \beta^\sharp)||_{\mathrm{op}}\}$*, where* a_λ *(resp.* a_μ*) is the eigenvalue of the Casimir operator of an irreducible complex representation belonging to* λ *(resp.* μ*) with respect to* $cB_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}}$ *(*$c = \frac{3}{4}$ *in case of* (I) *and* $c = \frac{1}{2}$ *in case of* (II)*),* V_{ρ_λ} *is the representation space of an irreducible representation* ρ_λ *belonging to* λ*,* m_λ *is the dimension of* V_{ρ_λ}, $\mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}})$ *is the* L*-module of all* L*-homomorphisms from* V_{ρ_λ} *to* $(\mathfrak{m}^\perp)^{\mathbf{c}}$ *and* $\beta^\sharp$ *is the element of* $\mathfrak{b}$ *defined by* $\beta(\cdot) = B_{\mathfrak{g}}(\beta^\sharp, \cdot)$*,* $\hat{c} := \frac{1}{3}$ *in case of* (I) *and* $\hat{c} := 1$ *in case of* (II)*, and* $||\cdot||_{\mathrm{op}}$ *is the operator norm of* $(\cdot)$ *with respect to* $B_{\mathfrak{g}}|_{\mathfrak{b}\times\mathfrak{b}}$.

In the last section, we give examples of a Hermann action $H \curvearrowright G/K$ and $Z_0 \in \widetilde{C}$ as in Theorem A and estimate the indices of $M := H(\mathrm{Exp}\, Z_0)$ for some of the examples by using Theorem A.

2 Basic notions and facts

In this section, we recall some basic notions and facts.

Jacobi operators

Let $f : (M, g) \hookrightarrow (\widetilde{M}, \widetilde{g})$ be a minimal isometric immersion of a compact Riemannian manifold (M, g) into another Riemannian manifold $(\widetilde{M}, \widetilde{g})$. Denote by $T^\perp M$ the normal bundle of $f(M)$ and $\Gamma(T^\perp M)$ the space of all normal vector fields of $f(M)$. Also, denote by ∇(resp. $\nabla^\perp$) the Levi-Civita connection of g (resp. the normal connection of $f(M)$) and A the shape tensor of $f(M)$. Let f_t $(-\varepsilon < t < \varepsilon)$ be a C^∞-family of immersions of M into $\widetilde{M}$ with $f_0 = f$, where ε is a positive number. Define a map $F : M \times (-\varepsilon, \varepsilon) \to \widetilde{M}$ by $F(x, t) := f_t(x)$ $((x, t) \in M \times (-\varepsilon, \varepsilon))$. Denote by $\mathrm{Vol}(M, f_t^*\widetilde{g})$ the volume of $(M, f_t^*\widetilde{g})$ and dv the volume element of g, where $f_t^*\widetilde{g}$ is the metric induced form $\widetilde{g}$ by f_t. Then we have the following second variational formula:

$$\frac{d^2}{dt^2}|_{t=0}\mathrm{Vol}(M, f_t^*\widetilde{g}) = \int_M \widetilde{g}(\mathcal{J}(F_*(\frac{\partial}{\partial t}|_{t=0})_\perp), F_*(\frac{\partial}{\partial t}|_{t=0})_\perp) dv.$$

Here F_* is the differential of F, $(\cdot)_\perp$ is the normal component of $(\cdot)$, $\mathcal{J}$ is a so-called Jacobi operator defined by $\mathcal{J} := -\triangle^\perp + \mathcal{R} - \mathcal{A}\,(: \Gamma(T^\perp M) \to \Gamma(T^\perp M))$, where $\triangle^\perp$ is the rough Laplacian operator defined by ∇ and $\nabla^\perp$, $\mathcal{A}$ is defined by $g(\mathcal{A}(v), w) = \mathrm{Tr}(A_v \circ A_w)$ $(v, w \in \Gamma(T^\perp M))$ and $\mathcal{R}$ is defined by $g(\mathcal{R}(v), w) = -\mathrm{Tr}(R(\cdot, v)w)$ $(v, w \in \Gamma(T^\perp M))$. Set $E^\perp_\lambda := \{v \in \Gamma(T^\perp M) \,|\, \mathcal{J}(v) = \lambda v\}$ for each $\lambda \in \mathbb{R}$. The dimension of $\sum\limits_{\lambda < 0} E^\perp_\lambda$ (resp. $E^\perp_0$) is called the *index* (resp. *nullity*) of f (or M).

The eigenvalues of the Casimir operators

Let H be a simple complex Lie group and $D(H)$ be the set of all equivalence classes of irreducible (complex) representations of H. Fix a Cartan subalgebra $\widetilde{\mathfrak{a}}$ of the Lie algebra $\mathfrak{h}$ of H. Let $\triangle$ be the root system of $\mathfrak{h}$ with respect to $\widetilde{\mathfrak{a}}$, $\triangle_+$ the positive root system of $\triangle$ under some lexicographic ordering of the dual space $\widetilde{\mathfrak{a}}^*$ of $\widetilde{\mathfrak{a}}$ and $\Pi = \{\alpha_1, \cdots, \alpha_r\}$ be a simple root system of $\triangle_+$. Fix a $\mathrm{Ad}(H)$-invariant non-degenerate $\mathbb{C}$-bilinear form $\langle\ ,\ \rangle$ of $\mathfrak{h}$. Define $\Lambda_i \in \widetilde{\mathfrak{a}}^*$ $(i = 1, \cdots, r)$ by $\dfrac{2\langle \alpha_j, \Lambda_i\rangle}{\langle \alpha_j, \alpha_j\rangle} = \delta_{ij}$ $(1 \le i, j \le r)$. A one-to-one correspondence between $D(H)$ and $\mathbb{Z}_+\{\Lambda_1, \cdots, \Lambda_r\}$ $(:= \{\sum_{i=1}^r z_i\Lambda_i \,|\, z_i \in \mathbb{Z}_+\})$ is obtained by assigning the highest weight of ρ to each $\lambda = [\rho] \in D(H)$, where $[\rho]$ is the equivalence class of a representation ρ of H. We denote the equivalence class of the irreducible representation of H corresponding to $\sum\limits_{i=1}^{r} z_i\Lambda_i$ by $(z_1, \cdots, z_r)$. The Casimir operator C_ρ of an irreducible representation ρ of H with respect to $\langle\ ,\ \rangle$ is defined by $C_\rho := \sum\limits_{i=1}^{m} \rho_{*e}(e_i)^2$, where $(e_1, \cdots, e_m)$ is an orthonormal base of $\mathfrak{h}$ with respect to $\langle\ ,\ \rangle$ and e is the identity element of H. By Schur's lemma, we have $C_\rho = a_\rho \mathrm{id}_{\mathfrak{h}}$ for some $a_\rho \in \mathbb{R}$, where $\mathrm{id}_{\mathfrak{h}}$ is the identity transformation of $\mathfrak{h}$. According to the Freudenthal's formula, we have

$$a_\rho = -\langle \Lambda, \Lambda + \sum_{\alpha \in \triangle_+} \alpha \rangle, \tag{2.1}$$

where Λ is the highest weight of ρ.

The canonical connection

Let H/L be a reductive homogeneous space and $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$ be a reductive decomposition (i.e., $[\mathfrak{h}, \mathfrak{m}] \subset \mathfrak{m}$), where $\mathfrak{h}$ (resp. $\mathfrak{l}$) is the Lie algebra of H (resp. L) and $\pi : P \to H/L$ be a principal G-bundle, where G is a Lie group. Assume that H acts on P and that $\pi(h \cdot u) = h \cdot \pi(u)$ for any $u \in P$ and $h \in H$. Then there uniquely exists a connection ω of P such that, for any $X \in \mathfrak{m}$ and any $u \in P$, $t \mapsto (\exp tX)(u)$ is a horizontal curve with respect to ω, where exp is the exponential map of H. This connection ω is called the *canonical connection* of P with respect to the reductive decomposition $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$.

The rough Laplacian operator with respect to the canonical connection

Let H be a Lie group and H/L be a reductive homogeneous space with a reductive decomposition $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$, where $\mathfrak{h}$ is the Lie algebra of H. The subspace $\mathfrak{m}$ is identified with $T_{eL}(H/L)$. Let B be an $\mathrm{Ad}(H)$-invariant inner product of $\mathfrak{h}$ such that $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$ is an orthogonal decomposition with respect to B. Denote by $\langle\ ,\ \rangle$ the H-invariant metric on H/L induced from $B|_{\mathfrak{m}\times\mathfrak{m}}$ and ∇ the Levi-Civita connection of $\langle\ ,\ \rangle$. Let $\pi : H \to H/L$ be the natural projection, $\sigma : L \to GL(W)$

be a unitary representation of L and $E_\sigma := H \times_{\sigma(L)} W$ be the associated complex vector bundle of the L-bundle $\pi : H \to H/L$ with respect to σ. The Lie group H acts on H and H/L naturally. Also, each $h(\in H)$ gives a linear isomorphism of W onto the fibre $(E_\sigma)_{\pi(h)}$. Denote by $\Gamma(E_\sigma)$ the space of all sections of E_σ and set $C^\infty(H,W)_\sigma := \{f \in C^\infty(H,W) \,|\, f(hl) = \sigma(l^{-1})f(h)\,(\forall\, h \in H, \forall\, l \in L)\}$, where $C^\infty(H,W)$ is the space of all W-valued C^∞-functions on H. Define a map $\Psi : \Gamma(E_\sigma) \to C^\infty(H,W)_\sigma$ by $\Psi(\xi)(h) = h^{-1}\cdot \xi_{\pi(h)}$ ($\xi \in \Gamma(E_\sigma)$, $h \in H$). This map Ψ is a linear isomorphism preserving the H-action. Take an orthonormal base $(e_1, \cdots, e_m)$ of $\mathfrak{h}$ with respect to B with $e_i \in \mathfrak{l}$ ($i = 1, \cdots, n$) and $e_b \in \mathfrak{m}$ ($b = n+1, \cdots, m$), where $n := \dim \mathfrak{l}$. Let $\mathcal{C}_H$ (: $C^\infty(H,W) \to C^\infty(H,W)$) be the Casimir differential operator of H with respect to B, that is, $\mathcal{C}_H(f) = \sum\limits_{i=1}^{m} \widetilde{e_i}(\widetilde{e_i}f)$, where $\widetilde{e_i}$ is the left-invariant vector field induced from e_i. Also, let $\mathcal{C}_\sigma$ be the Casimir operator of σ with respect to $B|_{\mathfrak{l}\times\mathfrak{l}}$. For $f \in C^\infty(H,W)_\sigma$, we can show $\mathcal{C}_H(f) = \mathcal{C}_\sigma \circ f + \sum\limits_{b=n+1}^{m} \widetilde{e_b}(\widetilde{e_b}f)$. Let ∇^ω be the connection of E_σ induced from the canonical connection ω of $\pi : H \to H/L$ with respect to the reductive decomposition $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$ and $\triangle^{E_\sigma}$ be the rough Laplacian operator of E_σ with respect to ∇^ω and ∇. Set $\widetilde{\triangle^{E_\sigma}} := \Psi \circ \triangle^{E_\sigma} \circ \Psi^{-1}$. Then we have $\widetilde{\triangle^{E_\sigma}} f = \sum\limits_{b=n+1}^{m} \widetilde{e_b}(\widetilde{e_b}f)$ ($f \in C^\infty(H,W)_\sigma$) by Proposition 2.3 of [Oh]. Furthermore, by Corollary 2.5 of [O], we have the following relation.

Lemma 2.1([O]). *For each $f \in C^\infty(H,W)_\sigma$, we have*

$$\widetilde{\triangle^{E_\sigma}} f = \mathcal{C}_H(f) - \mathcal{C}_\sigma \circ f.$$

Shape operators of orbits of Hermann actions

Let $H \curvearrowright G/K$ be a Hermann action and θ (resp. τ) be an involution of G with $(\mathrm{Fix}\,\theta)_0 \subset K \subset \mathrm{Fix}\,\theta$ (resp. $(\mathrm{Fix}\,\tau)_0 \subset H \subset \mathrm{Fix}\,\tau$). Assume that $\theta \circ \tau = \tau \circ \theta$. Let $\mathfrak{f}$, $\mathfrak{p}$, $\mathfrak{h}$, $\mathfrak{q}$, $\mathfrak{b}$, $\mathfrak{p}_\beta$, $\triangle'$, $\triangle'^V_+$, $\triangle'^H_+$, $\widetilde{C}$ and C be as in Introduction. Take a simplex σ of $\partial\widetilde{C}$ and $Z_0 \in \sigma$. Set $M := H(\mathrm{Exp}\,Z_0)$ and $g_0 := \exp\,Z_0$, where Exp is the exponential map of G/K at eK and exp is the exponential map of G. Denote by A the shape tensor of M. The tangent space $T_{\mathrm{Exp}\,Z_0}M$ of M at $\mathrm{Exp}\,Z_0$ is given by

$$T_{\mathrm{Exp}\,Z_0}M = g_{0*}\left(\mathfrak{z}_{\mathfrak{p}\cap\mathfrak{h}}(\mathfrak{b}) + \sum_{\beta\in\triangle'^V_+\setminus\triangle'^V_{Z_0}}(\mathfrak{p}_\beta\cap\mathfrak{q}) + \sum_{\beta\in\triangle'^H_+\setminus\triangle'^H_{Z_0}}(\mathfrak{p}_\beta\cap\mathfrak{h})\right) \tag{2.2}$$

and hence

$$T^{\perp}_{\mathrm{Exp}\,Z_0}M = g_{0*}\left(\mathfrak{b} + \sum_{\beta\in{\triangle'}^V_{Z_0}}(\mathfrak{p}_\beta\cap\mathfrak{q}) + \sum_{\beta\in{\triangle'}^H_{Z_0}}(\mathfrak{p}_\beta\cap\mathfrak{h})\right), \tag{2.3}$$

where ${\triangle'}^V_{Z_0}$ and ${\triangle'}^H_{Z_0}$ are as stated in Introduction. Take $v\in\mathfrak{b}$. The slice representation $\rho^S_{Z_0}(h) = h_{*\mathrm{Exp}\,Z_0}|_{T^{\perp}_{\mathrm{Exp}\,Z_0}M}$ $(h\in H_{Z_0})$. Then we have $\underset{h\in H_{Z_0}}{\cup}\rho^S_{Z_0}(h)(\mathfrak{b}) = T^{\perp}_{\mathrm{Exp}\,Z_0}M$ and

$$\begin{aligned}
&A_{\rho^S_{Z_0}(h)(v)}|_{\rho^S_{Z_0}(h)(g_{0*}(\mathfrak{z}_{\mathfrak{p}\cap\mathfrak{h}}(\mathfrak{b})))} = 0,\\
&A_{\rho^S_{Z_0}(h)(v)}|_{\rho^S_{Z_0}(h)(g_{0*}(\mathfrak{p}_\beta\cap\mathfrak{q}))} = -\frac{\beta(v)}{\tan\beta(Z_0)}\mathrm{id}\ \ (\beta\in{\triangle'}^V_+\setminus{\triangle'}^V_{Z_0}),\\
&A_{\rho^S_{Z_0}(h)(v)}|_{\rho^S_{Z_0}(h)(g_{0*}(\mathfrak{p}_\beta\cap\mathfrak{h}))} = \beta(v)\tan\beta(Z_0)\mathrm{id}\ \ (\beta\in{\triangle'}^H_+\setminus{\triangle'}^H_{Z_0}),
\end{aligned} \tag{2.4}$$

where $h\in H_{Z_0}$.

3 Proof of Theorem A

In this section, we shall prove Theorem A. Let $H \curvearrowright G/K$ and $M = H(\mathrm{Exp}\,Z_0)$ be as in Theorem A. We use the notations in Introduction.

Proof of Theorem A. Since Z_0 satisfies the condition (I) or (II) in Theorem A, it follows from (2.4) that M is a (non-totally geodesic) minimal submanifold. In the sequel, we shall show that the estimate inequality in Theorem A holds. First we note that $\mathfrak{a} = \mathfrak{b}$, that is, $\triangle_+ = \triangle'_+$ because the cohomogeneity of the H-action is equal to the rank of G/K. We suffice to show in the case where the G-invariant metric induced from $B_{\mathfrak{g}}|_{\mathfrak{p}\times\mathfrak{p}}$ is given to G/K. Denote by $\langle\ ,\ \rangle$ this G-invariant metric and g_I the metric on M induced from $\langle\ ,\ \rangle$. For simplicity, set $L := H_{Z_0}$ and let $\mathfrak{l}$ be the Lie algebra of L.
(Step I) In this step, we shall show that there exists a reductive decomposition $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$ satisfying $B_{\mathfrak{g}}(\mathfrak{l},\mathfrak{m}) = 0$. Easily we have

$$\mathfrak{l} = \mathfrak{z}_{\mathfrak{f}\cap\mathfrak{h}}(\mathfrak{b}) + \sum_{\beta\in{\triangle'}^V_{Z_0}}(\mathfrak{f}_\beta\cap\mathfrak{h}) + \sum_{\beta\in{\triangle'}^H_{Z_0}}(\mathfrak{p}_\beta\cap\mathfrak{h}). \tag{3.1}$$

Define a subspace $\mathfrak{m}$ of $\mathfrak{h}$ by

$$\mathfrak{m} := \mathfrak{z}_{\mathfrak{p}\cap\mathfrak{h}}(\mathfrak{b}) + \sum_{\beta\in{\triangle'}^V_+\setminus{\triangle'}^V_{Z_0}}(\mathfrak{f}_\beta\cap\mathfrak{h}) + \sum_{\beta\in{\triangle'}^H_+\setminus{\triangle'}^H_{Z_0}}(\mathfrak{p}_\beta\cap\mathfrak{h}).$$

Easily we can show that $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$ is a reductive decomposition and that $B_{\mathfrak{g}}(\mathfrak{l}, \mathfrak{m}) = 0$.
(Step II) Define a diffeomorphism $\psi : H/L \to M$ by $\psi(hL) := h \cdot \mathrm{Exp}\, Z_0$ $(h \in H)$. In this step, we shall show that $(\psi^* g_I)_{eL} = cB_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}}$, where c is as in the statement of Theorem A. For each $X \in \mathfrak{m}(= T_{eL}(H/L) = T_{\mathrm{Exp}(Z_0)}M)$, denote by X^* the Killing field on M associated with X, that is, $X^*_p := \frac{d}{dt}|_{t=0}(\exp tX)(p)\,(p \in M)$. From the definition of ψ, we have $\psi_{*eL}X = X^*_{\mathrm{Exp}\, Z_0}$. Take $S_\beta \in \mathfrak{f}_\beta \cap \mathfrak{h}$ and let T_β be the element of $\mathfrak{p}_\beta \cap \mathfrak{h}$ such that $\mathrm{ad}(b)(S_\beta) = \beta(b) T_\beta$ for any $b \in \mathfrak{b}$, where $\beta \in (\triangle'^V_+ \setminus \triangle'^V_{Z_0}) \cup (\triangle'^H_+ \setminus \triangle'^H_{Z_0})$. Then we have

$$\psi_{*eL}(S_\beta) = (S^*_\beta)_{\mathrm{Exp}\, Z_0} = -\sin\beta(Z_0)(\exp\, Z_0)_*(T_\beta) \tag{3.2}$$

and

$$\psi_{*eL}(T_\beta) = (T^*_\beta)_{\mathrm{Exp}\, Z_0} = \cos\beta(Z_0)(\exp\, Z_0)_*(T_\beta). \tag{3.3}$$

Hence, if Z_0 is as in (I), then we have $(\psi^* g_I)_{eL}(S_\beta, S_\beta) = \frac{3}{4}B_{\mathfrak{g}}(S_\beta, S_\beta)$ and $(\psi^* g_I)_{eL}$ $(T_\beta, T_\beta) = \frac{3}{4}B_{\mathfrak{g}}(T_\beta, T_\beta)$. Therefore we obtain $(\psi^* g_I)_{eL} = \frac{3}{4}B_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}}$. Also, if Z_0 is as in (II), then we have $(\psi^* g_I)_{eL}(S_\beta, S_\beta) = \frac{1}{2}B_{\mathfrak{g}}(S_\beta, S_\beta)$ and $(\psi^* g_I)_{eL}(T_\beta, T_\beta) = \frac{1}{2}B_{\mathfrak{g}}(T_\beta, T_\beta)$. Therefore we obtain $(\psi^* g_I)_{eL} = \frac{1}{2}B_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}}$.
(Step III) Let ω be the canonical connection of the principal L-bundle $\pi : H \to H/L(= M)$ with respect to the reductive decomposition $\mathfrak{h} = \mathfrak{l} + \mathfrak{m}$ and $F^\perp(M)$ be the normal frame bundle of M. Define a map $\eta : H \to F^\perp(M)$ by $\eta(h) = h_* u_0$ $(h \in H)$, where u_0 is an arbitrary fixed element of $F^\perp(M)_{\mathrm{Exp}\, Z_0}$, where $F^\perp(M)_{\mathrm{Exp}\, Z_0}$ is the fibre of $F^\perp(M)$ over $\mathrm{Exp}\, Z_0$. This map η is an embedding. By identifying H with $\eta(H)$, we regard $\pi : H \to H/L(= M)$ as a subbundle of $F^\perp(M)$. Denote by the same symbol ω the connection of $F^\perp(M)$ induced from ω and $\nabla^{\perp\omega}$ the linear connection on $T^\perp M$ associated with ω. Denote by $\nabla^\perp$ the normal connection of the submanifold M. In this step, we shall show that $\nabla^{\perp\omega} = \nabla^\perp$. Take $v \in \mathfrak{b}\,(\subset T^\perp_{\mathrm{Exp}\, Z_0}M)$. Let $Z : [0,1] \to \mathfrak{b}$ be a C^∞-curve with $Z(0) = Z_0$ and $Z((0,1]) \subset \widetilde{C}$. Set $M_s := H(\mathrm{Exp}\, Z(s))$ $(0 \leq s \leq 1)$. Denote by A^s the shape tensor of M_s and $\widetilde{\nabla}$ the Levi-Civita connection of G/K. Let $\widetilde{v}^s$ be the H-equivariant normal vector field of M_s $(0 < s \leq 1)$ arising from v. Since M_s $(0 < s \leq 1)$ is a principal orbit of the Hermann action H, $\widetilde{v}^s$ is well-defined and it is a parallel normal vector field. Take $X \in \mathfrak{f}_\beta \cap \mathfrak{h}\,(\subset \mathfrak{m})$ $(\beta \in \triangle'^V_+ \setminus \triangle'^V_{Z_0})$. Then, by using (2.4), we have

$$\widetilde{\nabla}_{X^*_{\mathrm{Exp}\, Z(s)}} \widetilde{v}^s = -A^s_v X^*_{\mathrm{Exp}\, Z(s)} = \frac{\beta(v)}{\tan\beta(Z_0)} X^*_{\mathrm{Exp}\, Z(s)}.$$

and hence

$$\widetilde{\nabla}_{X^*_{\mathrm{Exp}\, Z_0}} (\exp\, tX)_{*\mathrm{Exp}(Z_0)}(v) = \lim_{s\to+0} \widetilde{\nabla}_{X^*_{\mathrm{Exp}\, Z(s)}} \widetilde{v}^s = \frac{\beta(v)}{\tan\beta(Z_0)} X^*_{\mathrm{Exp}\, Z_0} \in T_{\mathrm{Exp}\, Z_0}M.$$

Hence we obtain $\nabla^{\perp}_{X^*_{\mathrm{Exp}\,Z_0}}(\exp tX)_{*\mathrm{Exp}(Z_0)}(v) = 0$. Take $Y \in \mathfrak{p}_\beta \cap \mathfrak{h}\,(\subset \mathfrak{m})$ $(\beta \in \triangle'^H_+ \setminus \triangle'^H_{Z_0})$. Then, by using (2.4), we have

$$\widetilde{\nabla}_{Y^*_{\mathrm{Exp}\,Z(s)}}\widetilde{v}^s = -A^s_v Y^*_{\mathrm{Exp}\,Z(s)} = -\beta(v)\tan\beta(Z_0)Y^*_{\mathrm{Exp}\,Z(s)}.$$

and hence

$$\begin{aligned}\widetilde{\nabla}_{Y^*_{\mathrm{Exp}\,Z_0}}(\exp tY)_{*\mathrm{Exp}\,Z_0}(v) &= \lim_{s\to+0}\widetilde{\nabla}_{Y^*_{\mathrm{Exp}\,Z(s)}}\widetilde{v}^s \\ &= -\beta(v)\tan\beta(Z_0)Y^*_{\mathrm{Exp}\,Z_0} \in T_{\mathrm{Exp}\,Z_0}M.\end{aligned}$$

Hence we obtain $\nabla^{\perp}_{Y^*_{\mathrm{Exp}\,Z_0}}(\exp tY)_{*\mathrm{Exp}(Z_0)}(v) = 0$. Therefore, it follows from the arbitrariness of X, Y and β that $t \mapsto (\exp t\hat{X})_{*\mathrm{Exp}\,Z_0}(v)$ is $\nabla^{\perp}$-parallel along $t \mapsto (\exp t\hat{X})(\mathrm{Exp}\,Z_0)$ for any $\hat{X} \in \mathfrak{m}$. Take any $h \in L$. Similarly we can show that $t \mapsto (\exp t\hat{X})_{*\mathrm{Exp}\,Z_0}(\rho^S_{Z_0}(h)(v))$ is $\nabla^{\perp}$-parallel along $t \mapsto (\exp t\hat{X})(\mathrm{Exp}\,Z_0)$ for any $\hat{X} \in \mathfrak{m}$, where $\rho^S_{Z_0}$ is the slice representation of the H-action at $\mathrm{Exp}\,Z_0$. Note that this fact has been showed in [IST] in different method. On the other hand, it follows from the definition of ω that $t \mapsto (\exp t\hat{X})_{*\mathrm{Exp}\,Z_0}(\rho^S_{Z_0}(h)(v))$ is $\nabla^{\perp\omega}$-parallel along $t \mapsto (\exp t\hat{X})(\mathrm{Exp}\,Z_0)$ for any $\hat{X} \in \mathfrak{m}$. Therefore we obtain $\nabla^{\perp} = \nabla^{\perp\omega}$.

(Final step) Now we shall prove Theorem A by using the facts in Steps I $\sim$ III. The space $\mathfrak{m}^{\perp} := (\exp Z_0)^{-1}_*(T^{\perp}_{\mathrm{Exp}\,Z_0}M)$ is regarded as the representation space of $\rho^S_{Z_0}$ under the identification of $\mathfrak{m}^{\perp}$ and $T^{\perp}_{\mathrm{Exp}\,Z_0}M$. Denote by $E_{\rho^S_{Z_0}}$ the associated vector bundle $H \times_{\rho^S_{Z_0}} \mathfrak{m}^{\perp}$ of the principal L-bundle $\pi : H \to H/L$ with respect to $\rho^S_{Z_0}$. This vector bundle $E_{\rho^S_{Z_0}}$ is identified with $T^{\perp}M$ under the correspondence $h \cdot v \leftrightarrow h_*((\exp Z_0)_*v)$ $(h \in H,\ v \in \mathfrak{m}^{\perp})$. Let $\Psi : \Gamma(E_{\rho^S_{Z_0}}) \to C^{\infty}(H, \mathfrak{m}^{\perp})_{\rho^S_{Z_0}}$ be a diffeomorphism stated in the previous section. Since $\psi^* g_I$ is an H-invariant metric induced from $cB_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}}$ and $\nabla^{\perp} = \nabla^{\perp\omega}$, it follows from Lemma 2.1 that the rough Laplacian operator $\triangle^{\perp}$ of $E_{\rho^S_{Z_0}}$ with respect to $\nabla^{\perp}$ and ∇ satisfies

$$(3.4) \qquad (\Psi \circ \triangle^{\perp} \circ \Psi^{-1})(f) = \mathcal{C}_H(f) - \mathcal{C}_{\rho^S_{Z_0}} \circ f \quad (f \in C^{\infty}(H, \mathfrak{m}^{\perp})_{\rho^S_{Z_0}}),$$

where $\mathcal{C}_H$ is the Casimir differential operator of H with respect to $cB_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}}$ and $\mathcal{C}_{\rho^S_{Z_0}}$ is the Casimir operator of $\rho^S_{Z_0}$ with respect to $cB_{\mathfrak{g}}|_{\mathfrak{l}\times\mathfrak{l}}$. Let $\rho_{G/H}$ be the isotropy representation of G/H, $\mathcal{C}_{\rho_{G/H}}$ the Casimir operator of $\rho_{G/H}$ with respect to $cB|_{\mathfrak{h}\times\mathfrak{h}}$ and μ the equivalence class of the complexification of $\rho_{G/H}$. The slice representation $\rho^S_{Z_0}$ is equivalent to the representation $(\rho_{G/H}|_L)|_{\mathfrak{m}^{\perp}}$ of L defined by $((\rho_{G/H}|_L)|_{\mathfrak{m}^{\perp}})(l) := \rho_{G/H}(l)|_{\mathfrak{m}^{\perp}}$ $(l \in L)$. Hence, from (3.4) and the definition of $\mathcal{R}$, we have

$$(3.5) \qquad (\Psi \circ (\triangle^{\perp} - \mathcal{R}) \circ \Psi^{-1})(f) = \mathcal{C}_H(f) - \mathcal{C}_{\rho_{G/H}} \circ f \quad (f \in C^{\infty}(H, \mathfrak{m}^{\perp})_{\rho^S_{Z_0}}).$$

Since G/H is irreducible or $G/H = G'/H' \times G'/H'$ for some irreducible symmetric space G'/H', we have $\mathcal{C}_{\rho_{G/H}} = a_\mu \mathrm{id}$ for some constant a_μ. Hence we have

$$(\Psi \circ (\triangle^\perp - \mathcal{R}) \circ \Psi^{-1})(f) = \mathcal{C}_H(f) - a_\mu f \quad (f \in C^\infty(H, \mathfrak{m}^\perp)_{\rho^S_{Z_0}}). \tag{3.6}$$

Let $\lambda(= [\rho_\lambda]) \in D(H)$ and V_{ρ_λ} be the representaion space of ρ_λ. Define a map $\eta_{\rho_\lambda} : V_{\rho_\lambda} \otimes \mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}}) \to C^\infty(H, (\mathfrak{m}^\perp)^{\mathbf{c}})_{(\rho^S_{Z_0})^{\mathbf{c}}}$ by $(\eta_{\rho_\lambda}(v \otimes \phi))(h) := \phi(\rho_\lambda(h^{-1})(v))$ $(v \in V_{\rho_\lambda},\ \phi \in \mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}}),\ h \in H)$, where $\mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}})$ is the space of all L-homomorphisms from V_{ρ_λ} to $(\mathfrak{m}^\perp)^{\mathbf{c}}$. This map η_{ρ_λ} is injective. Set $\Gamma_\lambda((T^\perp M)^{\mathbf{c}}) := \Psi^{-1}(\eta_{\rho_\lambda}(V_{\rho_\lambda} \otimes \mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}})))$. Then it is shown that $\sum_{\lambda \in D(H)} \Gamma_\lambda((T^\perp M)^{\mathbf{c}})$ (direct sum) is uniformly dense in $\Gamma((T^\perp M)^{\mathbf{c}})$ with respect to the uniformly topology. Also, it follows from (3.6) that

$$(\triangle^\perp - \mathcal{R})^{\mathbf{c}}|_{\Gamma_\lambda((T^\perp M)^{\mathbf{c}})} = (a_\lambda - a_\mu)\mathrm{id}. \tag{3.7}$$

On the other hand, it follows from (2.4), $\tan^2 \beta(Z_0) = \frac{1}{\hat{c}}$ $(\beta \in {\triangle'}^V_+ \setminus {\triangle'}^V_{Z_0})$ and $\tan^2 \beta(Z_0) = \hat{c}$ $(\beta \in {\triangle'}^H_+ \setminus {\triangle'}^H_{Z_0})$ that

$$0 < \langle \mathcal{A}(v), v \rangle \leq \Big|\Big| \sum_{\beta \in \triangle'_+ \setminus ({\triangle'}^V_{Z_0} \cup {\triangle'}^H_{Z_0})} m_\beta \hat{c} (\beta \otimes \beta^\sharp) \Big|\Big|_{\mathrm{op}} \langle v, v \rangle \tag{3.8}$$

for any $v(\neq 0) \in \Gamma(T^\perp M)$. From (3.7) and (3.8), we have

$$\begin{aligned} &\left(a_\mu - a_\lambda - \Big|\Big| \sum_{\beta \in \triangle'_+ \setminus ({\triangle'}^V_{Z_0} \cup {\triangle'}^H_{Z_0})} m_\beta \hat{c} (\beta \otimes \beta^\sharp) \Big|\Big|_{\mathrm{op}} \right) \int_M \langle v, v \rangle dv \\ &\leq \int_M \langle \mathcal{J}(v), v \rangle dv \ < \ (a_\mu - a_\lambda) \int_M \langle v, v \rangle dv \\ &\qquad\qquad\qquad\qquad (v \in \Gamma_\lambda((T^\perp M)^{\mathbf{c}}) \cap \Gamma(T^\perp M)). \end{aligned} \tag{3.9}$$

Therefore the index $i(M)$ is estimated as in the statement of Theorem A. In particular, from (3.9), we have

$$\int_M \langle \mathcal{J}(v), v \rangle dv \ < \ 0 \qquad (v \in \Gamma_\mu((T^\perp M)^{\mathbf{c}}) \cap \Gamma(T^\perp M)).$$

Therefore M is unstable. This completes the proof. q.e.d.

Remark 3.1.

(i) The normal vector field $\xi^{v \otimes \phi}$ is explicitly described as

$$(\xi^{v \otimes \phi})_{h \mathrm{Exp}\, Z_0} = h_*((\exp\, Z_0)_*(\rho_{G/H}(h^{-1})(v))) \quad (h \in H). \tag{3.10}$$

(ii) The closure $\overline{\widetilde{C}}$ of $\widetilde{C}$ stated in Introduction is a simplicial complex. If $M := H\mathrm{Exp}\,Z_0$ ($Z_0 \in \overline{\widetilde{C}}$) is totally geodesic, then we have $Z_0 = 0$ in the case where G/K is other than a sphere. Then, for each element $\xi^{v\otimes\phi} := \Psi^{-1}(\eta_\mu(v \otimes \phi))$ ($v \in V_{\rho_\mu}$, $\phi \in \mathrm{Hom}_L(V_{\rho_\mu}, (\mathfrak{m}^\perp)^{\mathbf{c}})$) of $\Gamma_\mu((T^\perp M)^{\mathbf{c}}) \cap \Gamma(T^\perp M))$, there exists a Killing vector field X on G/K such that the normal component $(X|_M)_\perp$ of $X|_M$ is equal to $\xi^{v\otimes\phi}$. This normal vector field $\xi^{v\otimes\phi}$ and X are as in Fig. 1. Hence we have $\int_M \langle \mathcal{J}(\xi^{v\otimes\phi}), \xi^{v\otimes\phi}\rangle dv = 0$, which is derived from Theorem A because M is totally geodesic. If Z_0 satisfies the condition (I) or (II) in Introduction, then $M = H\mathrm{Exp}\,Z_0$ is not totally geodesic and each element $\xi^{v\otimes\phi}$ of $\Gamma_\mu((T^\perp M)^{\mathbf{c}}) \cap \Gamma(T^\perp M))$ is as in Fig. 2. There does not exist a Killing vector field X on G/K satisfying $(X|_M)_\perp = \xi^{v\otimes\phi}$. For example, in the case where $G/K = SO(3)/SO(2)$, $H = SO(2)$ and where M is a great circle in $SO(3)/SO(2)$, there exists a Killing vector field X on $SO(3)/SO(2)$ satisfying $(X|_M)_\perp = \xi^{v\otimes\phi}$ for any $v(\neq 0) \in \mathfrak{m}^\perp$ and the orthogonal projection ψ of $\mathfrak{q}$ onto $\mathfrak{m}^\perp$ (see Fig. 3). However, in the case where M is a small circle, there does not exist a Killing vector field X on $SO(3)/SO(2)$ satisfying $(X|_M)_\perp = \xi^{v\otimes\phi}$ for any $v(\neq 0) \in \mathfrak{m}^\perp$ and the orthogonal projection ψ of $\mathfrak{q}$ onto $\mathfrak{m}^\perp$ (see Fig. 4 and 5). Even if a small circle is not minimal, this fact indicates that, in the case where $M = H\mathrm{Exp}\,Z_0$ is minimal but not totally geodesic, there does not exist a Killing vector field X on G/K satisfying $(X|_M)_\perp = \xi^{v\otimes\phi}$ for each element $\xi^{v\otimes\phi}$ of $\Gamma_\mu((T^\perp M)^{\mathbf{c}}) \cap \Gamma(T^\perp M))$.


$Z_0 = 0$
$\overline{\widetilde{C}}$
$\mathfrak{b}$
in fact
$Z_0 = 0$
$\overline{\widetilde{C}}$
$\mathfrak{b}$
$M = H\mathrm{Exp}\,Z_0 = H(eK)$
Hp
$\mathrm{Exp}(\mathfrak{p} \cap \mathfrak{q})$
$\mathrm{Exp}(\overline{\widetilde{C}})$
$\mathrm{Exp}(\mathfrak{b})$
$\mathrm{Exp}\,Z_0 = eK$
p
's : $\xi^{v\otimes\phi}$
's : X

(X : the Killing vector field on G/K such that $(X|_M)^\perp = \xi^{v\otimes\phi}$)

Fig. 1.


$\overline{\widetilde{C}}$
Z_0
$\mathfrak{b}$
in fact
$\overline{\widetilde{C}}$
Z_0
$\mathfrak{b}$
$M = H\mathrm{Exp}\,Z_0$
$\mathrm{Exp}(\mathfrak{p}\cap\mathfrak{q})$
$\mathrm{Exp}(\mathfrak{b})$
$\mathrm{Exp}(\overline{\widetilde{C}})$
$\mathrm{Exp}\,Z_0$
’s : $\xi^{v\otimes\phi}$

There does not exist a Killing vector field X on G/K s.t. $(X|_M)_\perp = \xi^{v\otimes\phi}$.

Fig. 2.


$(\xi^{v\otimes\phi})_{e^{\frac{\pi i}{2}}\mathrm{Exp}\,Z_0}(=0)$
$(\xi^{v\otimes\phi})_{e^{\frac{3\pi i}{4}}\mathrm{Exp}\,Z_0}$
$(\xi^{v\otimes\phi})_{e^{\frac{\pi i}{4}}\mathrm{Exp}\,Z_0}$
$M := SO(2)(\mathrm{Exp}\,Z_0)$
$(\xi^{v\otimes\phi})_{e^{\pi i}\mathrm{Exp}\,Z_0}$
$\mathrm{Exp}\,Z_0$
$\rho_\mu(e^{-\frac{\pi i}{4}})(v)$
v
$(\xi^{v\otimes\phi})_{\mathrm{Exp}\,Z_0}$
$\phi(\rho_\mu(e^{-\frac{\pi i}{4}})(v))$
$(\xi^{v\otimes\phi})_{e^{\frac{-3\pi i}{4}}\mathrm{Exp}\,Z_0}$
$(\xi^{v\otimes\phi})_{e^{\frac{-\pi i}{4}}\mathrm{Exp}\,Z_0}$
$eSO(2)$
$(\xi^{v\otimes\phi})_{e^{-\frac{\pi i}{2}}\mathrm{Exp}\,Z_0}(=0)$

$\xi^{v\otimes\phi}$ extends to a Killing vector field on $SO(3)/SO(2)$.

Fig. 3.


$(\xi^{v\otimes\phi})_{e^{\frac{\pi i}{2}}\operatorname{Exp} Z_0}(=0)$
$M := SO(2)(\operatorname{Exp} Z_0)$
(small circle)
$(\xi^{v\otimes\phi})_{e^{\frac{3\pi i}{4}}\operatorname{Exp} Z_0}$
$(\xi^{v\otimes\phi})_{e^{\pi i}\operatorname{Exp} Z_0}$
$(\xi^{v\otimes\phi})_{e^{\frac{\pi i}{4}}\operatorname{Exp} Z_0}$
$\rho_\mu(e^{-\frac{\pi i}{4}})(v)$
v
$\operatorname{Exp} Z_0$
$\phi(\rho_\mu(e^{-\frac{\pi i}{4}})(v))$
$(\xi^{v\otimes\phi})_{\operatorname{Exp} Z_0}$
$(\xi^{v\otimes\phi})_{e^{\frac{-3\pi i}{4}}\operatorname{Exp} Z_0}$
$eSO(2)$
$(\xi^{v\otimes\phi})_{e^{\frac{-\pi i}{4}}\operatorname{Exp} Z_0}$
$(\xi^{v\otimes\phi})_{e^{-\frac{\pi i}{2}}\operatorname{Exp} Z_0}(=0)$

There does not exist a Killing vector field X on $SO(3)/SO(2)$ s.t. $(X|_M)_\perp = \xi^{v\otimes\phi}$.

Fig. 4.


$SO(2)(\operatorname{Exp} Z_1)$ (small circle)
$SO(2)(\operatorname{Exp} Z_0)$
(great circle)
$(\xi^{v\otimes\phi})_{e^{\frac{\pi i}{4}}\operatorname{Exp} Z_0} = (X_\perp)_{e^{\frac{\pi i}{4}}\operatorname{Exp} Z_0}$
$(\xi^{v\otimes\phi})_{e^{\frac{\pi i}{4}}\operatorname{Exp} Z_1} \neq (X_\perp)_{e^{\frac{\pi i}{4}}\operatorname{Exp} Z_1}$
's : a Killing vector field

Fig. 5.

4 Examples

In this section, we give examples of a Hermann action $H \curvearrowright G/K$ and $Z_0 \in \widetilde{C}$ as in Theorem A and estimate the indices of $M := H(\operatorname{Exp} Z_0)$ for some of the examples by using Theorem A. We use the notations in Introduction.

Example 1. We consider the isotropy action of $SU(3n+3)/SO(3n+3)$. Then we have $\triangle_+ = \triangle'_+ = \triangle'^V_+$, which is of $(\mathfrak{a}_{3n+2})$-type. Let $\Pi = \{\beta_1, \cdots, \beta_{3n+2}\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 4, $\triangle'_+ = \{\beta_i + \cdots + \beta_j \,|\, 1 \leq i, j \leq 3n+2\}$ and that $m_\beta = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be the point of the closure of $\widetilde{C}$ defined by $\beta_{n+1}(Z_0) = \beta_{2n+2}(Z_0) = \frac{\pi}{3}$ and $\beta_i(Z_0) = 0$ ($i \in \{1, \cdots, 3n+2\} \setminus \{n+1, 2n+2\}$). This point Z_0 satisfies the condition (I) in Introduction.



Fig. 4.

Example 2. We consider the isotropy action of $SU(6n+6)/Sp(3n+3)$. Then we have $\triangle_+ = \triangle'_+ = \triangle'^V_+$, which is of $(\mathfrak{a}_{3n+2})$-type. Let $\Pi = \{\beta_1, \cdots, \beta_{3n+2}\}$ be a simple root system of $\triangle'_+$. We have $m_\beta = 4$ for any $\beta \in \triangle'_+$. Let Z_0 be the point of the closure of $\widetilde{C}$ defined by $\beta_{n+1}(Z_0) = \beta_{2n+2}(Z_0) = \frac{\pi}{3}$ and $\beta_i(Z_0) = 0$ ($i \in \{1, \cdots, 3n+2\} \setminus \{n+1, 2n+2\}$). This point Z_0 satisfies the condition (I) in Introduction.

Example 3. We consider the isotropy action of $SU(3)/S(U(1)\times U(2))$ (2-dimensional complex projective space). Then we have $\triangle_+ = \triangle'_+ = \triangle'^V_+ = \{\beta, 2\beta\}$, $m_\beta = 2$ and $m_{2\beta} = 1$. Let Z_0 be the point of $\widetilde{C}$ defined by $\beta(Z_0) = \frac{\pi}{3}$. This point Z_0 satisfies the condition (I) in Introduction.

Example 4. We consider the isotropy action of $Sp(3n+2)/U(3n+2)$. Then we have $\triangle_+ = \triangle'_+ = \triangle'^V_+$, which is of $(\mathfrak{c}_{3n+2})$-type. Let $\Pi = \{\beta_1, \cdots, \beta_{3n+2}\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 5. We have $m_\beta = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be the point of the closure of $\widetilde{C}$ defined by $\beta_{n+1}(Z_0) = \beta_{3n+2}(Z_0) = \frac{\pi}{3}$ and $\beta_i(Z_0) = 0$ ($i \in \{1, \cdots, 3n+2\} \setminus \{n+1, 3n+2\}$). This point Z_0 satisfies the condition (I) in Introduction.



Fig. 5.

Example 5. We consider the Hermann action $SO(2n) \curvearrowright SU(2n)/Sp(n)$. The cohomogeneity of this action is equal to the rank of $SU(2n)/Sp(n)$. Also we have $\triangle_+ = \triangle'_+ = \triangle'^V_+ = \triangle'^H_+$, which is of $(\mathfrak{a}_{n-1})$-type. Let $\Pi = \{\beta_1, \cdots, \beta_{n-1}\}$ be a simple root system of $\triangle'_+$. We have $m^V_\beta = m^H_\beta = 2$ for any $\beta \in \triangle'_+$. Let Z_0 be a point of the closure of $\widetilde{C}$ such that one or two of $\beta_1(Z_0), \cdots, \beta_{n-1}(Z_0)$ are equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_0)$ are equal to zero. This point Z_0 satisfies the condition (II) in Introduction.

Example 6. We consider the Hermann action $SO(n) \times SO(n) \curvearrowright (SU(n) \times SU(n))/SU(n)$. The cohomogeneity of this action is equal to the rank of $(SU(n) \times SU(n))/SU(n)$. Also we have $\triangle_+ = \triangle'_+ = \triangle'^V_+ = \triangle'^H_+$, which is of $(\mathfrak{a}_{n-1})$-type. Let $\Pi = \{\beta_1, \cdots, \beta_{n-1}\}$ be a simple root system of $\triangle'_+$. We have $m^V_\beta = m^H_\beta = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be a point of the closure of $\widetilde{C}$ such that one or two of $\beta_1(Z_0), \cdots, \beta_{n-1}(Z_0)$ are equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_0)$ are equal to zero. This point Z_0 satisfies the condition (II) in Introduction.

Example 7. We consider the Hermann action $U(n) \times U(n) \curvearrowright (Sp(n) \times Sp(n))/Sp(n)$. The cohomogeneity of this action is equal to the rank of $(Sp(n) \times Sp(n))/Sp(n)$. Also we have $\triangle_+ = \triangle'_+ = \triangle'^V_+ = \triangle'^H_+$, which is of $(\mathfrak{c}_n)$-type. Let $\Pi = \{\beta_1, \cdots, \beta_n\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 6. We have $m^V_\beta = m^H_\beta = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be a point of the closure of $\widetilde{C}$ such that one of $\beta_1(Z_0), \cdots, \beta_{n-1}(Z_0)$ is equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_0)$ are equal to zero. This point Z_0 satisfies the condition (II) in Introduction.



Fig. 6.

Example 8. We consider the Hermann action $Sp(4) \times Sp(4) \curvearrowright (E_6 \times E_6)/E_6$. 00The cohomogeneity of this action is equal to the rank of $(E_6 \times E_6)/E_6$. Also we have $\triangle_+ = \triangle'_+ = \triangle'^V_+ = \triangle'^H_+$, which is of $(\mathfrak{e}_6)$-type. Let $\Pi = \{\beta_1, \cdots, \beta_6\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 7. We have $m^V_\beta = m^H_\beta = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be a point of the closure of $\widetilde{C}$ such that one of $\beta_1(Z_0), \beta_2(Z_0), \beta_4(Z_0), \beta_5(Z_0), \beta_6(Z_0)$ is equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_0)$ are equal to zero. Also, let Z_1 be a point of the closure of $\widetilde{C}$ such that $\beta_1(Z_1)$ and $\beta_6(Z_1)$ are equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_1)$ are equal to zero. These points Z_0 and Z_1 satisfy the condition (II) in Introduction.



Fig. 7.

Example 9. We consider the Hermann action $SU(8) \times SU(8) \curvearrowright (E_7 \times E_7)/E_7$. The cohomogeneity of this action is equal to the rank of $(E_7 \times E_7)/E_7$. Also we have $\triangle_+ = \triangle'_+ = {\triangle'}^V_+ = {\triangle'}^H_+$, which is of $(\mathfrak{e}_7)$-type. Let $\Pi = \{\beta_1, \cdots, \beta_7\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 8. We have $m^V_\beta = m^H_\beta = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be a point of the closure of $\widetilde{C}$ such that one of $\beta_1(Z_0), \beta_2(Z_0), \beta_6(Z_0), \beta_7(Z_0)$ is equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_0)$ are equal to zero. This point Z_0 satisfies the condition (II) in Introduction.



Fig. 8.

Example 10. We consider the Hermann action $SO(16) \times SO(16) \curvearrowright (E_8 \times E_8)/E_8$. The cohomogeneity of this action is equal to the rank of $(E_8 \times E_8)/E_8$. Also we have $\triangle_+ = \triangle'_+ = {\triangle'}^V_+ = {\triangle'}^H_+$, which is of $(\mathfrak{e}_8)$-type. Let $\Pi = \{\beta_1, \cdots, \beta_8\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 9. 1We have $m^V_\beta = m^H_\beta = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be a point of the closure of $\widetilde{C}$ such that one of $\beta_1(Z_0), \beta_8(Z_0)$ is equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_0)$ are equal to zero. This point Z_0 satisfies the condition (II) in Introduction.



Fig. 9

Example 11. We consider the Hermann action $(Sp(3) \cdot SU(2)) \times (Sp(3) \cdot SU(2)) \curvearrowright (F_4 \times F_4)/F_4$. The cohomogeneity of this action is equal to the rank of $(F_4 \times F_4)/F_4$. Also we have $\triangle_+ = \triangle'_+ = {\triangle'}^V_+ = {\triangle'}^H_+$, which is of $(\mathfrak{f}_4)$-type. Let $\Pi = \{\beta_1, \cdots, \beta_4\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 10. We have $m_\beta^V = m_\beta^H = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be a point of the closure of $\widetilde{C}$ such that one of $\beta_1(Z_0), \beta_4(Z_0)$ is equal to $\frac{\pi}{4}$ and that other $\beta_i(Z_0)$ are equal to zero. This point Z_0 satisfies the condition (II) in Introduction.



Fig. 10.

Example 12. We consider the Hermann action $SU(2)^4 \curvearrowright (G_2 \times G_2)/G_2$. The cohomogeneity of this action is equal to the rank of $(G_2 \times G_2)/G_2$. Also we have $\triangle_+ = \triangle'_+ = \triangle'^V_+ = \triangle'^H_+$, which is of $(\mathfrak{g}_2)$-type. Let $\Pi = \{\beta_1, \beta_2\}$ be a simple root system of $\triangle'_+$, where we note that the Dynkin diagram of $\triangle'_+$ is as in Fig. 11. We have $m_\beta^V = m_\beta^H = 1$ for any $\beta \in \triangle'_+$. Let Z_0 be the point of the closure of $\widetilde{C}$ satisfying $\beta_2(Z_0) = \frac{\pi}{4}$ and $\beta_1(Z_0) = 0$. This point Z_0 satisfies the condition (II) in Introduction.



Fig. 11.

Now we give a recipi to estimate concretely the index of the minimal orbit $M = H(\operatorname{Exp} Z_0)$ for a Hermann action $H \curvearrowright G/K$ and $Z_0 \in \widetilde{C}$ in Theorem A.

Recipi. Denote by $\hat{B}$ the $\mathrm{Ad}(H)$-invariant inner product $cB_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}}$ of $\mathfrak{h}$, where c is as in Theorem A. Let $\mathfrak{h} = \mathfrak{h}_1 + \cdots + \mathfrak{h}_k$ be the irreducible decomposition of $\mathfrak{h}$. Denote by $\bar{B}_i$ $(i = 1, \cdots, k)$ the $\mathrm{Ad}(H_i)$-invariant inner product of $\mathfrak{h}_i$ satisfying $\max\limits_{\alpha\in\triangle^{\mathfrak{h}_i^{\mathbf{c}}}} \bar{B}_i(\bar{\alpha}^\sharp, \bar{\alpha}^\sharp) = 2$, where $H_i := \exp \mathfrak{h}_i$ and $\triangle^{\mathfrak{h}_i^{\mathbf{c}}}$ is the root system of $\mathfrak{h}_i^{\mathbf{c}}$ with respect to the complexification $\mathfrak{t}_i^{\mathbf{c}}$ of a maximal abelian subalgebra $\mathfrak{t}_i$ of $\mathfrak{h}_i$ and $\bar{\alpha}^\sharp$ is the element of $\mathfrak{t}_i$ defined by $\alpha|_{\mathfrak{t}_i}(\cdot) = \bar{B}_i(\bar{\alpha}^\sharp, \cdot)$. The complexification $\bar{B}_i^{\mathbf{c}}$ of $\bar{B}_i$ is the scalar product of $\mathfrak{h}_i^{\mathbf{c}}$ considered in [MP] (see Page 3 of [MP]). By Schur's lemma, $\hat{B}|_{\mathfrak{h}_i\times\mathfrak{h}_i} = c_i\bar{B}_i$ holds for some positive constant c_i and, for each $\lambda = [\rho_\lambda] = [\rho_{\lambda_1}\text{-}\cdots-\rho_{\lambda_k}] \in D(H)$ $(\rho_{\lambda_i} \in D(H_i))$, $a_{\lambda_i} = \frac{1}{c_i}\bar{a}_{\lambda_i}$ holds between the eigenvalues a_{λ_i} and $\bar{a}_{\lambda_i}$ of the Casimir operators of ρ_{λ_i} with respect to $\hat{B}|_{\mathfrak{h}_i\times\mathfrak{h}_i}$ and $\bar{B}_i$, where $\rho_{\lambda_1}-\cdots-\rho_{\lambda_k}$ is the representation of $H_1 \times \cdots \times H_k$ defined by $(\rho_{\lambda_1} - \cdots - \rho_{\lambda_k})(h_1, \cdots, h_k)(v_1 \otimes \cdots \otimes v_k) := \rho_{\lambda_1}(h_1)(v_1) \otimes \cdots \otimes \rho_{\lambda_k}(h_k)(v_k)$ $(h_i \in H_i,\ v_i \in V_{\rho_{\lambda_i}})$ (the representation space of $\rho_{\lambda_1} - \cdots - \rho_{\lambda_k}$ is $V_{\rho_{\lambda_1}} \otimes \cdots \otimes V_{\rho_{\lambda_k}}$). Hence the eigenvalue a_λ of the Casimir operator ρ_λ with respect to $\hat{B}$ is equal to $\sum\limits_{i=1}^{k} a_{\lambda_i} = \sum\limits_{i=1}^{k} \frac{1}{c_i}\bar{a}_{\lambda_i}$.

(Step I) We calculate these constants $c_1, \cdots, c_k$.

Let $\Pi = \{\beta_1, \cdots, \beta_r\}$ be the simple root system of $\triangle'_+$ and θ_{ij} $(1 \leq i, j \leq r)$ the angle between $\beta_i^\sharp$ and $\beta_j^\sharp$, where $\beta_i^\sharp$ is the element of $\mathfrak{b}$ defined by $\beta_i(\cdot) = B_{\mathfrak{g}}(\beta_i^\sharp, \cdot)$. Set $l_i := \sqrt{B_{\mathfrak{g}}(\beta_i^\sharp, \beta_i^\sharp)}$ $(i = 1, \cdots, r)$. For each $\beta \in \triangle'_+$, we set $\beta = \sum\limits_{i=1}^{r} n_\beta^i \beta_i$. For each $\mathbf{v} \in \mathfrak{b}$, we set $\mathbf{v} = \sum\limits_{i=1}^{r} v^i \beta_i^\sharp$. Then we have $B_{\mathfrak{g}}(\mathbf{v}, \mathbf{v}) = \sum\limits_{1 \leq i,j \leq r} v^i v^j l_i l_j \cos\theta_{ij}$ and

$$(4.3) \qquad \begin{aligned} & || \sum_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} m_\beta \hat{c} (\beta \otimes \beta^\sharp) ||_{\mathrm{op}} \\ & = \max_{v \in \mathfrak{b} \setminus \{0\}} \left(\sum_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} m_\beta \hat{c} \beta(v)^2 \right) / B_{\mathfrak{g}}(v, v) \\ & = \max_{(v^1, \cdots, v^r) \in \mathbf{R}^r \setminus \{(0, \cdots, 0)\}} \left(\sum_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} (m_\beta \hat{c} \right. \\ & \left. \times \sum_{1 \leq i,j,k,u \leq r} n_\beta^i n_\beta^k v^j v^u l_i l_j l_k l_u \cos\theta_{ij} \cos\theta_{ku}) \right) / \left(\sum_{1 \leq i,j \leq r} v^i v^j l_i l_j \cos\theta_{ij} \right). \end{aligned}$$

Let C_1 and C_2 be the symmetric matrices of degree r such that

$${}^t\mathbf{v} C_1 \mathbf{v} = \sum_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} (m_\beta \hat{c} \sum_{1 \leq i,j,k,u \leq r} n_\beta^i n_\beta^k v^j v^u l_i l_j l_k l_u \cos\theta_{ij} \cos\theta_{ku})$$

and

$${}^t\mathbf{v} C_2 \mathbf{v} = \sum_{1 \leq i,j \leq r} v^i v^j l_i l_j \cos\theta_{ij}$$

for all $\mathbf{v} = (v^1, \cdots, v^r) \in \mathbb{R}^r$. To calculate $|| \sum\limits_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} m_\beta \hat{c} (\beta \otimes \beta^\sharp) ||_{\mathrm{op}}$ $= \max\limits_{\mathbf{v} \in \mathbf{R}^r \setminus \{(0, \cdots, 0)\}} \dfrac{{}^t\mathbf{v} C_1 \mathbf{v}}{{}^t\mathbf{v} C_2 \mathbf{v}}$, we have only to calculate the maximum of positive numbers κ's such that ${}^t\mathbf{v}(C_1 - \kappa C_2)\mathbf{v} = 0$ for some $\mathbf{v} \neq (0, \cdots 0)$, That is, we have only to calculate the maximum of κ's such that $C_1 - \kappa C_2$ is neither a positive operator nor a negative operator.

(Step II) We calculate C_1 and C_2, and estimate the maximum of κ's satisfying the above condition strictly.

(Step III) We calculate $\bar{a}_{\lambda_i}$ (as in (Step I)) in terms of Table 1 of $I_0(\lambda_i)(= \dim V_{\rho_{\lambda_i}})$

and $I_2(\lambda_i)(=-\frac{\mathrm{rank}\,\mathfrak{h}_i\cdot I_0(\lambda_i)\cdot\bar{a}_{\lambda_i}}{\dim\mathfrak{h}_i})$ in [MP], where we take λ_i as Λ in [MP].
As its result, we can calculate a_λ's ($\lambda\in D(H)$) and estimate $a_\mu-||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}}$ strictly.

(Step IV) By comparing the values of a_μ and $a_\mu-||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}}$ with those of a_λ's ($\lambda\in D(H)$), we list up the elements of $D_1(H)$ and $D_2(H)$, where μ is the equivalence class of the complexification of the isotropy representation of G/H.

(Step V) By using the relation (3.1), we analyze the Lie algebra $\mathfrak{l}$ of the isotropy group $L:=H_{Z_0}$.

(Step VI) For each element $\lambda=[\rho_\lambda]$ of $D_1(H)$ and $D_2(H)$, we investigate how $\rho_\lambda|_L$ is decomposed irreducibly by using Table 2 (the branching rule) in [MP].
Also we investigate how $(\rho_{G/H}|_L)^{\mathbf{c}}|_{(\mathfrak{m}^\perp)^{\mathbf{c}}}$ is decomposed irreducibly.

(Step VII) By using the irreducible decompositions of $(\rho_{G/H}|_L)^{\mathbf{c}}|_{(\mathfrak{m}^\perp)^{\mathbf{c}}}$ and $\rho_\lambda|_L$'s ($\lambda\in D_2(H)$), we calculate $\dim\mathrm{Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}})$ and estimate the index $i(M)$ of M by using the estimate inequality in Theorem A.

By using this recipi, we shall estimate the indices of some of the minimal orbits $M=H(\mathrm{Exp}\,Z_0)$ in Examples 1 $\sim$ 12. First we consider the case where $G/K=SU(6)/SO(6)$, $H=SO(6)$, and $M=SO(6)(\mathrm{Exp}\,Z_0)$ ($\beta_2(Z_0)=\beta_4(Z_0)=\frac{\pi}{3}$, $\beta_1(Z_0)=\beta_3(Z_0)=\beta_5(Z_0)=0$) (the case of $n=1$ in Example 1). Then we have $\triangle'=\triangle=\triangle^{\mathfrak{g}^{\mathbf{c}}}$ (($\mathfrak{a}_5$)-type). Since $B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}}=3B_{\mathfrak{h}}$ (see Pages 187, 188 of [Hel]), we have $(\psi^*g_I)_{eL}=cB_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}}=\frac{3}{4}(B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}})|_{\mathfrak{m}\times\mathfrak{m}}=\frac{9}{4}B_{\mathfrak{h}}|_{\mathfrak{m}\times\mathfrak{m}}$, which implies $\hat{B}=\frac{9}{4}B_{\mathfrak{h}}$. Let $\mathfrak{t}$ be a maximal abelian subalgebra of $\mathfrak{h}$ and $\triangle^{\mathfrak{h}^{\mathbf{c}}}$ be the root system of $\mathfrak{h}^{\mathbf{c}}$ with respect to $\mathfrak{t}^{\mathbf{c}}$ and define $\alpha^\sharp\,(\in\mathfrak{t})$ (resp. $\hat{\alpha}^\sharp\,(\in\mathfrak{t})$) by $\alpha(\cdot)=B_{\mathfrak{h}}(\alpha^\sharp,\cdot)=\hat{B}(\hat{\alpha}^\sharp,\cdot)$. Take $\alpha\in\triangle^{\mathfrak{h}^{\mathbf{c}}}$. Since $B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp)=\frac{1}{4}$ (see Page 188 of [Hel]) and $\hat{B}=\frac{9}{4}B_{\mathfrak{h}}$, we have $\hat{B}(\hat{\alpha}^\sharp,\hat{\alpha}^\sharp)=\frac{4}{9}B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp)=\frac{1}{9}$. From this relation, it follows that $\hat{B}=18\bar{B}$. Hence we have $a_\lambda=\frac{1}{18}\bar{a}_\lambda$ for each $\lambda\in D(H)$. Since $\triangle'$ is ($\mathfrak{a}_5$)-type and $\Pi=\{\beta_1,\cdots,\beta_5\}$ is a simple root system of $\triangle'_+$, we have $\triangle'_+=\{\beta_i+\cdots+\beta_j\,|\,1\leq i,j\leq 5\}$, $l_i=\sqrt{B_{\mathfrak{g}}(\beta_i^\sharp,\beta_i^\sharp)}=\frac{1}{\sqrt{6}}$ ($i=1,\cdots,5$), $\theta_{i,i+1}=\frac{2}{3}\pi$ ($i=1,\cdots,4$) and $\theta_{ij}=\frac{\pi}{2}$ ($j-i\geq 2$). Hence the symmetric matrices C_1 and C_2 in (Step II) are given by

$$C_1=\frac{1}{216}\begin{pmatrix}-4 & 8 & -4 & 0 & 0\\ 8 & -12 & 12 & -5 & 0\\ -4 & 12 & -16 & 12 & -4\\ 0 & -5 & 12 & -12 & 8\\ 0 & 0 & -4 & 8 & -4\end{pmatrix}$$

and

$$C_2=\frac{1}{12}\begin{pmatrix} 2 & -1 & 0 & 0 & 0\\ -1 & 2 & -1 & 0 & 0\\ 0 & -1 & 2 & -1 & 0\\ 0 & 0 & -1 & 2 & -1\\ 0 & 0 & 0 & -1 & 2\end{pmatrix},$$

respectively. It is shown that the maximum of the value κ such that $C_1-\kappa C_2$ is neither a positive operator nor a negative operator is bigger than $\frac{19}{72}$ and smaller than $\frac{5}{18}$. Thus we have $\frac{19}{72}<||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})}m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\rm op}<\frac{5}{18}$. Since the equivalence class μ of the complexification of the isotropy representation of G/H is $(0\,2\,0)$, we have $\bar{a}_\mu=-12$ in terms of Table 1 in [MP] and the relation (2.3) in Page 10 of [MP]. Hence we have $a_\mu=-\frac{2}{3}$. Therefore we have $-\frac{17}{18}<a_\mu-||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})}m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\rm op}<-\frac{67}{72}$. Similarly we can calculate a_λ for other $\lambda\in D(H)$. By comparing the values of a_μ and $a_\mu-||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})}m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\rm op}$ with those of a_λ's ($\lambda\in D(H)$ s.t. $\lambda\neq\mu$), we have $D_1(H)=\{(0\,0\,0),\,(0\,0\,1),\,(0\,1\,0),\,(0\,1\,1),\,(0\,0\,2),\,(1\,0\,1),(0\,2\,0)\}$ and $D_2(H)=\{(0\,0\,0),\,(0\,0\,1),\,(0\,1\,0),\,(0\,1\,1),\,(0\,0\,2),\,(1\,0\,1),(0\,2\,0),\,(0\,0\,3),\,(1\,0\,2),\,(0\,1\,2),\,(1\,1\,1)\}$. Since $\triangle'_{Z_0}=\{\beta_1,\beta_3,\beta_5\}$ and $\mathfrak{z}_{\mathfrak{h}}(\mathfrak{b})=\{0\}$, we have $\mathfrak{l}=\mathfrak{h}_{\beta_1}+\mathfrak{h}_{\beta_3}+\mathfrak{h}_{\beta_5}$, where $\mathfrak{h}_{\beta_i}=\{X\in\mathfrak{h}\,|\,{\rm ad}(b)^2(X)=-\beta_i(b)^2X\ (\forall\,b\in\mathfrak{b})\}$ $(i=1,3,5)$. Thus $\mathfrak{l}$ is abelian, that is, $\mathfrak{l}=\mathfrak{so}(2)+\mathfrak{so}(2)+\mathfrak{so}(2)$. Hence we have $\dim{\rm Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}})=\dim V_{\rho_\lambda}\cdot\dim(\mathfrak{m}^\perp)^{\mathbf{c}}$ for each $\lambda\in D(H)$. Also we have $\dim\mathfrak{m}^\perp=8$. Therefore we have

$$\sum_{\lambda\in D_1(H)}m_\lambda\dim{\rm Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}})=8\sum_{\lambda\in D_1(H)}m_\lambda^2=9424$$

and

$$\sum_{\lambda\in D_2(H)}m_\lambda\dim{\rm Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}})=8\sum_{\lambda\in D_2(H)}m_\lambda^2=71960.$$

See Table 1 of [MP] about m_μ and m_λ's. Therefore we obtain the following result.

Proposition 4.1. *Let $\Pi=\{\beta_1,\cdots,\beta_5\}$ be the simple root system of the positive root system $\triangle_+$ of $SU(6)/SO(6)$ ($\underset{\beta_1}{\circ}-\underset{\beta_2}{\circ}-\underset{\beta_3}{\circ}-\underset{\beta_4}{\circ}-\underset{\beta_5}{\circ}$) and Z_0 be the element of $\mathfrak{b}$ with $\beta_2(Z_0)=\beta_4(Z_0)=\frac{\pi}{3}$ and $\beta_1(Z_0)=\beta_3(Z_0)=\beta_5(Z_0)=0$. Then the orbit $M:=SO(6)({\rm Exp}(Z_0))$ of the isotropy action of $SU(6)/SO(6)$ is a (non-totally geodesic) unstable minimal submanifold and we have $9424\leq i(M)\leq 71960$.*

Next we consider the case where $G/K=SU(9)/SO(9)$, $H=SO(9)$, and $M=SO(9)({\rm Exp}\,Z_0)$ ($\beta_3(Z_0)=\beta_6(Z_0)=\frac{\pi}{3}$, $\beta_i(Z_0)=0\,(i\neq 3,6)$) (the case of $n=2$ in

Example 1). Then we have $\triangle' = \triangle = \triangle^{\mathfrak{g}^{\mathbf{c}}}$ (($\mathfrak{a}_8$)-type). Since $B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}} = \frac{18}{7}B_{\mathfrak{h}}$ (see Page 187, 189 of [Hel]), we have $(\psi^* g_I)_{eL} = cB_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}} = \frac{3}{4}(B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}})|_{\mathfrak{m}\times\mathfrak{m}} = \frac{27}{14}B_{\mathfrak{h}}|_{\mathfrak{m}\times\mathfrak{m}}$, which implies $\hat{B} = \frac{27}{14}B_{\mathfrak{h}}$. Let $\mathfrak{t}$ be a maximal abelian subalgebra of $\mathfrak{h}$ and $\triangle^{\mathfrak{h}^{\mathbf{c}}}$ be the root system of $\mathfrak{h}^{\mathbf{c}}$ with respect to $\mathfrak{t}^{\mathbf{c}}$ and define $\alpha^\sharp\,(\in \mathfrak{t})$ (resp. $\hat{\alpha}^\sharp\,(\in \mathfrak{t})$) by $\alpha(\cdot) = B_{\mathfrak{h}}(\alpha^\sharp, \cdot) = \hat{B}(\hat{\alpha}^\sharp, \cdot)$. Take $\alpha \in \triangle^{\mathfrak{h}^{\mathbf{c}}}$. Since $B_{\mathfrak{h}}(\alpha^\sharp, \alpha^\sharp) = \frac{1}{7}$ or $\frac{1}{14}$ (see Page 189 of [Hel]) and $\hat{B} = \frac{27}{14}B_{\mathfrak{h}}$, we have $\hat{B}(\hat{\alpha}^\sharp, \hat{\alpha}^\sharp) = \frac{14}{27}B_{\mathfrak{h}}(\alpha^\sharp, \alpha^\sharp) = \frac{1}{27}$ or $\frac{2}{27}$. From this relation, it follows that $\hat{B} = 27\bar{B}$. Hence we have $a_\lambda = \frac{1}{27}\bar{a}_\lambda$ for each $\lambda \in D(H)$. Since $\triangle'$ is ($\mathfrak{a}_8$)-type and $\Pi = \{\beta_1, \cdots, \beta_8\}$ is a simple root system of $\triangle'_+$, we have $\triangle'_+ = \{\beta_i + \cdots + \beta_j \,|\, 1 \leq i, j \leq 8\}$, $l_i = \sqrt{B_{\mathfrak{g}}(\beta_i^\sharp, \beta_i^\sharp)} = \frac{1}{3}$ $(i = 1, \cdots, 8)$, $\theta_{i,i+1} = \frac{2}{3}\pi$ $(i = 1, \cdots, 7)$ and $\theta_{ij} = \frac{\pi}{2}$ $(j - i \geq 2)$. Hence the symmetric matrices C_1 and C_2 in (Step II) are given by

$$C_1 = \frac{1}{972}\begin{pmatrix} 12 & -6 & 0 & 0 & 0 & 0 & 0 & 0 \\ -6 & 12 & -6 & 0 & 0 & 0 & 0 & 0 \\ 0 & -6 & 14 & -6 & 0 & -1 & 0 & 0 \\ 0 & 0 & -6 & 12 & -6 & 0 & 0 & 0 \\ 0 & 0 & 0 & -6 & 12 & -6 & 0 & 0 \\ 0 & 0 & -1 & 0 & -6 & 14 & -6 & 0 \\ 0 & 0 & 0 & 0 & 0 & -6 & 12 & -6 \\ 0 & 0 & 0 & 0 & 0 & 0 & -6 & 12 \end{pmatrix}$$

and

$$C_2 = \frac{1}{18}\begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix},$$

respectively. It is shown that the maximum of the value κ such that $C_1 - \kappa C_2$ is neither a positive operator nor a negative operator is bigger than $\frac{4}{27}$ and smaller than $\frac{1}{6}$. Thus we have $\frac{4}{27} < ||\sum\limits_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} m_\beta \hat{c}(\beta \otimes \beta^\sharp)||_{\mathrm{op}} < \frac{1}{6}$. Since the equivalence class μ of the complexification of the isotropy representation of G/H is $(2\,0\,0\,0)$, we have $\bar{a}_\mu = -18$ in terms of Table 1 in [MP] and the relation (2.3) in Page 10 of [MP]. Hence we have $a_\mu = -\frac{2}{3}$. Therefore we have $-\frac{5}{6} < a_\mu - ||\sum\limits_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} m_\beta \hat{c}(\beta \otimes \beta^\sharp)||_{\mathrm{op}} < -\frac{22}{27}$. Similarly we can calculate a_λ for other $\lambda \in D(H)$. By comparing the values of a_μ and $a_\mu - ||\sum\limits_{\beta \in \triangle'_+ \setminus (\triangle'^V_{Z_0} \cup \triangle'^H_{Z_0})} m_\beta \hat{c}(\beta \otimes \beta^\sharp)||_{\mathrm{op}}$ with those of a_λ's

($\lambda \in D(H)$ s.t. $\lambda \neq \mu$), we have $D_1(H) = \{(0\,0\,0\,0),\ (1\,0\,0\,0),\ (0\,0\,0\,1), (0\,1\,0\,0), (2\,0\,0\,0),\ (0\,0\,1\,0), (1\,0\,0\,1)\}$ and $D_2(H) = \{(0\,0\,0\,0),\ (1\,0\,0\,0),\ (0\,0\,0\,1), (0\,1\,0\,0), (2\,0\,0\,0)\ (0\,0\,1\,0), (1\,0\,0\,1), (0\,0\,0\,2)\}$. Since $\triangle'_{Z_0} = \{\beta_1, \beta_2, \beta_1+\beta_2, \beta_4, \beta_5, \beta_4+\beta_5, \beta_7, \beta_8, \beta_7+\beta_8\}$ and $\mathfrak{z}_{\mathfrak{h}}(\mathfrak{b}) = \{0\}$, we have $\mathfrak{l} = \sum\limits_{i\in\{1,4,7\}} (\mathfrak{h}_{\beta_i} + \mathfrak{h}_{\beta_{i+1}} + \mathfrak{h}_{\beta_i+\beta_{i+1}})$. From this relation, $\mathfrak{l} = \mathfrak{so}(3) + \mathfrak{so}(3) + \mathfrak{so}(3)$ follows. Hence, by using Table 2 (the branching rules) in [MP], we have the following table:

λ	$\lambda\vert_L$	m_λ
$(0\,0\,0\,0)$	$(0-0-0)$	1
$(1\,0\,0\,0)$	$3(0-0-2)$	9
$(0\,0\,0\,1)$	$2(1-1-1)$	16
$(0\,1\,0\,0)$	$3(0-0-2) \oplus 3(0-2-2)$	36
$\mu = (2\,0\,0\,0)$	$2(0-0-0) \oplus 3(0-2-2) \oplus 3(0-0-4)$	44
$(0\,0\,1\,0)$	$3(0-0-0) \oplus 6(0-2-2) \oplus (2-2-2)$	84
$(0\,0\,0\,2)$	$6(0-0-2) \oplus 3(0-2-2) \oplus 3(2-2-2)$	126
$(1\,0\,0\,1)$	$4(1-1-1) \oplus 6(1-1-3)$	128

Table 1.

In this table, $(\lambda_1 - \lambda_2 - \lambda_3)$ ($\lambda_1, \lambda_2, \lambda_3 \in D(SO(3))$) is as stated in Recipi. Also, since $[\rho_\mu|_L] = 2(0-0-0) \oplus 3(0-2-2) \oplus 3(0-0-4)$ and $\dim \mathfrak{m}^\perp = 17$, we have

$$[(\rho_\mu|_L)|_{(\mathfrak{m}^\perp)^{\mathbf{c}}}] = 2(0-0-0) \oplus 3(0-0-4).$$

Hence, we have

$$\sum_{\lambda\in D_1(H)} m_\lambda \dim \mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}}) = \sum_{\lambda\in D_2(H)} m_\lambda \dim \mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}}) = 978.$$

Therefore we obtain the following result.

Proposition 4.2. *Let* $\Pi = \{\beta_1, \cdots, \beta_8\}$ *be the simple root system of the positive root system* $\triangle_+$ *of* $SU(9)/SO(9)$ ($\underset{\beta_1}{\circ} - \underset{\beta_2}{\circ} - \cdots - \underset{\beta_8}{\circ}$) *and* Z_0 *be the element of* $\mathfrak{b}$ *with* $\beta_3(Z_0) = \beta_6(Z_0) = \frac{\pi}{3}$ *and* $\beta_i(Z_0) = 0$ ($i \neq 3, 6$). *Then the orbit* $M := SO(9)(\mathrm{Exp}(Z_0))$ *of the isotropy action of* $SU(9)/SO(9)$ *is a (non-totally geodesic) unstable minimal submanifold and we have* $i(M) = 978$.

Next we consider the case where $G/K = SU(12)/Sp(6)$, $H = Sp(6)$, and $M = Sp(6)(\mathrm{Exp}\, Z_0)$ ($\beta_2(Z_0) = \beta_4(Z_0) = \frac{\pi}{3}$, $\beta_i(Z_0) = 0\,(i \neq 2, 4)$) (the case of $n = 1$ in Example 2) . Since the Satake diagram of G/K is as in Fig. 12 and $H = K$, we have $\triangle' = \triangle$ and the multiplicity of each root of $\triangle'$ is equal to 4.



$(\widetilde{\beta}_i|_{\mathfrak{b}} = \beta_i)$

Fig. 12.

Also, since $B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}} = \frac{12}{7}B_{\mathfrak{h}}$ (see Page 187, 190 of [Hel]), we have $(\psi^* g_I)_{eL} = cB_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}} = \frac{3}{4}(B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}})|_{\mathfrak{m}\times\mathfrak{m}} = \frac{9}{7}B_{\mathfrak{h}}|_{\mathfrak{m}\times\mathfrak{m}}$, which implies $\hat{B} = \frac{9}{7}B_{\mathfrak{h}}$. Let $\mathfrak{t}$ be a maximal abelian subalgebra of $\mathfrak{h}$ and $\triangle^{\mathfrak{h}^{\mathbf{c}}}$ be the root system of $\mathfrak{h}^{\mathbf{c}}$ with respect to $\mathfrak{t}^{\mathbf{c}}$ and define $\alpha^\sharp\,(\in\,\mathfrak{t})$ (resp. $\hat{\alpha}^\sharp\,(\in\,\mathfrak{t})$) by $\alpha(\cdot) = B_{\mathfrak{h}}(\alpha^\sharp,\cdot) = \hat{B}(\hat{\alpha}^\sharp,\cdot)$. Take $\alpha \in \triangle^{\mathfrak{h}^{\mathbf{c}}}$. Since $B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp) = \frac{1}{7}$ or $\frac{1}{14}$ (see Page 190 of [Hel]) and $\hat{B} = \frac{9}{7}B_{\mathfrak{h}}$, we have $\hat{B}(\hat{\alpha}^\sharp,\hat{\alpha}^\sharp) = \frac{7}{9}B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp) = \frac{1}{9}$ or $\frac{1}{18}$. From this relation, it follows that $\hat{B} = 18\bar{B}$. Hence we have $a_\lambda = \frac{1}{18}\bar{a}_\lambda$ for each $\lambda \in D(H)$. Since $\triangle'$ is $(\mathfrak{a}_5)$-type and $\Pi = \{\beta_1,\cdots,\beta_5\}$ is a simple root system of $\triangle'_+$, we have $\triangle'_+ = \{\beta_i+\cdots+\beta_j\,|\,1 \le i,j \le 5\}$, $l_i = \sqrt{B_{\mathfrak{g}}(\beta_i^\sharp,\beta_i^\sharp)} = \sqrt{\frac{1}{2}B_{\mathfrak{g}}(\widetilde{\beta}_i^\sharp,\widetilde{\beta}_i^\sharp)} = \frac{1}{2\sqrt{6}}$ $(i=1,\cdots,5)$, $\theta_{i,i+1} = \frac{2}{3}\pi$ $(i=1,\cdots,4)$ and $\theta_{ij} = \frac{\pi}{2}$ $(j-i \ge 2)$. Hence the symmetric matrices C_1 and C_2 in (Step II) are given by

$$C_1 = \frac{1}{1728}\begin{pmatrix} -8 & 16 & -8 & 0 & 0 \\ 16 & -24 & 24 & -10 & 0 \\ -8 & 24 & -32 & 24 & -8 \\ 0 & -10 & 24 & -24 & 16 \\ 0 & 0 & -8 & 16 & -8 \end{pmatrix}$$

and

$$C_2 = \frac{1}{48}\begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix},$$

respectively. It is shown that the maximum of the value κ such that $C_1 - \kappa C_2$ is neither a positive operator nor a negative operator is bigger than $\frac{19}{72}$ and smaller than $\frac{5}{18}$. Thus we have $\frac{19}{72} < ||\sum\limits_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}} < \frac{5}{18}$. Since the equivalence class μ of the complexification of the isotropy representation of G/H is $(0\,1\,0\,0\,0\,0)$, we have $\bar{a}_\mu = -12$ in terms of Table 1 in [MP] and the relation (2.3) in Page 10 of [MP]. Hence we have $a_\mu = -\frac{2}{3}$. Therefore we have $-\frac{17}{18} < a_\mu - ||\sum\limits_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}} < -\frac{67}{72}$. Similarly we can calculate a_λ for other $\lambda \in D(H)$. By comparing the values of a_μ and $a_\mu - ||\sum\limits_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}}$ with those of a_λ's ($\lambda \in D(H)$ s.t. $\lambda \ne \mu$), we have $D_1(H) = \{(0\,0\,0\,0\,0\,0),\ (1\,0\,0\,0\,0\,0),\ (0\,1\,0\,0\,0\,0)\}$ and $D_2(H) = \{(0\,0\,0\,0\,0\,0),\ (1\,0\,0\,0\,0\,0), (0\,1\,0\,0\,0\,0), (2\,0\,0\,0\,0\,0),\ (0\,0\,1\,0\,0\,0)\}$.

Since $\triangle'_{Z_0} = \{\beta_1, \beta_3, \beta_5\}$, we have $\mathfrak{l} = \mathfrak{z}_{\mathfrak{h}}(\mathfrak{b}) + \mathfrak{h}_{\beta_1} + \mathfrak{h}_{\beta_3} + \mathfrak{h}_{\beta_5}$. Also we have $\dim \mathfrak{l} = \dim G - (\dim G/K - \dim(\mathfrak{b} + \mathfrak{p}_{\beta_1} + \mathfrak{p}_{\beta_3} + \mathfrak{p}_{\beta_5}))$. From these facts, we have $\mathfrak{l} = 6\mathfrak{so}(2) + 3\mathfrak{so}(4) (= 6\mathfrak{so}(2) + 6\mathfrak{su}(2))$. Hence, by using Table 2 (the branching rules) in [MP], we have the following table:

λ	$\lambda\|_L$	m_λ
$(0\,0\,0\,0\,0\,0)$	$(0-0-0-0-0-0)$	1
$(1\,0\,0\,0\,0\,0)$	$6(0-0-0-0-0-1)$	12
$\mu = (0\,1\,0\,0\,0\,0)$	$5(0-0-0-0-0-0) \oplus 15(0-0-0-0-1-1)$	65
$(2\,0\,0\,0\,0\,0)$	$6(0-0-0-0-0-2) \oplus 15(0-0-0-0-1-1)$	78
$(0\,0\,1\,0\,0\,0)$	$9(0-0-0-0-0-0) \oplus 45(0-0-0-0-1-1)$ $\oplus 15(0-0-1-1-1-1)$	208

Table 2.

Also, since $[\rho_\mu|_L] = 5(0-0-0-0-0-0) \oplus 15(0-0-0-0-1-1)$, $\mathfrak{m}^\perp = \mathfrak{b} + \mathfrak{p}_{\beta_1} + \mathfrak{p}_{\beta_3} + \mathfrak{p}_{\beta_5}$, $\dim \mathfrak{b} = 5$ and $\dim \mathfrak{p}_{\beta_i} = 4$ $(i = 1, 3, 5)$, we have

$$[(\rho_\mu|_L)|_{(\mathfrak{m}^\perp)^{\mathbf{c}}}] = 5(0-0-0-0-0-0) \oplus 3(0-0-0-0-1-1).$$

From these facts, we have

$$\sum_{\lambda \in D_1(H)} m_\lambda \dim \mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}}) = 4555$$

and

$$\sum_{\lambda \in D_2(H)} m_\lambda \dim \mathrm{Hom}_L(V_{\rho_\lambda}, (\mathfrak{m}^\perp)^{\mathbf{c}}) = 45505.$$

Therefore we obtain the following result.

Proposition 4.3. *Let $\Pi = \{\beta_1, \cdots, \beta_5\}$ be the simple root system of the positive root system $\triangle_+$ of $SU(12)/Sp(6)$ ($\underset{\beta_1}{\circ} - \underset{\beta_2}{\circ} - \underset{\beta_3}{\circ} - \underset{\beta_4}{\circ} - \underset{\beta_5}{\circ}$) and Z_0 be the element of $\mathfrak{b}$ with $\beta_2(Z_0) = \beta_4(Z_0) = \frac{\pi}{3}$ and $\beta_1(Z_0) = \beta_3(Z_0) = \beta_5(Z_0) = 0$. Then the orbit $M := Sp(6)(\mathrm{Exp}(Z_0))$ of the isotropy action of $SU(12)/Sp(6)$ is a (non-totally geodesic) unstable minimal submanifold and we have $4555 \leq i(M) \leq 45505$.*

Next we consider the case where $G/K = SU(3)/S(U(1) \times U(2))$, $H = S(U(1) \times U(2))$, and $M = S(U(1) \times U(2))(\mathrm{Exp}\, Z_0)$ $(\beta(Z_0) = \frac{\pi}{3})$ (Example 3). Since the Satake diagram of G/K is $\underset{\widetilde{\beta}_1}{\circ} - \underset{\widetilde{\beta}_2}{\circ}$ and $H = K$, we have $\triangle' = \triangle = \{\beta, 2\beta\}$ and the multiplicity of β (resp. 2β) is equal to 2 (resp. 1), where $\widetilde{\beta}_i|_{\mathfrak{b}} = \beta$ $(i = 1, 2)$. Since $B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}} = \frac{3}{2} B_{\mathfrak{h}}$ (see Page 187 of [Hel]), we have $(\psi^* g_I)_{eL} = cB_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}} =$

$\frac{3}{4}(B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}})|_{\mathfrak{m}\times\mathfrak{m}} = \frac{9}{8}B_{\mathfrak{h}}|_{\mathfrak{m}\times\mathfrak{m}}$, which implies $\hat{B} = \frac{9}{8}B_{\mathfrak{h}}$. Let $\mathfrak{t}$ be a maximal abelian subalgebra of $\mathfrak{h}$ and $\triangle^{\mathfrak{h}^{\mathbf{c}}}$ be the root system of $\mathfrak{h}^{\mathbf{c}}$ with respect to $\mathfrak{t}^{\mathbf{c}}$ and define $\alpha^\sharp\,(\in\mathfrak{t})$ (resp. $\hat{\alpha}^\sharp\,(\in\mathfrak{t})$) by $\alpha(\cdot) = B_{\mathfrak{h}}(\alpha^\sharp,\cdot) = \hat{B}(\hat{\alpha}^\sharp,\cdot)$. Take $\alpha\in\triangle^{\mathfrak{h}^{\mathbf{c}}}$. Since $B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp) = \frac{1}{2}$ (see Page 187 of [Hel]) and $\hat{B} = \frac{9}{8}B_{\mathfrak{h}}$, we have $\hat{B}(\hat{\alpha}^\sharp,\hat{\alpha}^\sharp) = \frac{8}{9}B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp) = \frac{4}{9}$. From this relation, it follows that $\hat{B} = \frac{9}{2}\bar{B}$. Hence we have $a_\lambda = \frac{2}{9}\bar{a}_\lambda$ for each $\lambda\in D(H)$. Since $\triangle'_+ = \{\beta,2\beta\}$, we have $l_i = \sqrt{B_{\mathfrak{g}}(\beta^\sharp,\beta^\sharp)} = \frac{1}{2}\sqrt{B_{\mathfrak{g}}(\widetilde{\beta}_1^\sharp,\widetilde{\beta}_1^\sharp)} = \frac{1}{2\sqrt{3}}$. Hence the symmetric matrices C_1 and C_2 in (Step II) are given by $C_1 = (\frac{1}{72})$ and $C_2 = (\frac{1}{12})$, respectively. It is shown that the only value κ such that $C_1 - \kappa C_2$ is equal to $\frac{1}{6}$. This implies $||\sum\limits_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\rm op} = \frac{1}{6}$. Since the equivalence class μ of the complexification of the isotropy representation of G/H is $(0)\otimes(0)\otimes(3)$, we can show $\bar{a}_\mu = -\frac{15}{2}$ in terms of the fact related to irreducible representations of $\mathfrak{su}(2)$ stated in Page 13 of [MP] and the relation (2.3) in Page 10 of [MP], where we note that $\mathfrak{h} = \mathfrak{s}u(2) + \mathfrak{s}o(2)$. Hence we have $a_\mu = -\frac{5}{3}$. Therefore we have $a_\mu - ||\sum\limits_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\rm op} = -\frac{11}{6}$. Similarly we can calculate a_λ for other $\lambda\in D(H)$. By comparing the values of a_μ and $a_\mu - ||\sum\limits_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\rm op}$ with those of a_λ's ($\lambda\in D(H)$ s.t. $\lambda\neq\mu$), we have $D_1(H) = D_2(H) = \{(0),\,(1),\,(2),\,(3)\}$. On the other hand, we have $\mathfrak{l} = \mathfrak{z}_{\mathfrak{h}}(\mathfrak{b}) = \mathfrak{so}(2)$ and $\dim\mathfrak{m}^\perp = 1$. Hence we have $\dim{\rm Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}}) = m_\lambda\dim(\mathfrak{m}^\perp)^{\mathbf{c}} = m_\lambda$. Hence we have $\sum\limits_{\lambda\in D_1(H)} m_\lambda\dim{\rm Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}}) = \sum\limits_{\lambda\in D_2(H)} m_\lambda\dim{\rm Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}}) = 30$. Therefore we obtain the following result.

Proposition 4.4. *Let* $\triangle_+ := \{\beta,2\beta\}$ *be the positive root system of* $SU(3)/S(U(1)\times U(2))$ *and* Z_0 *be the element of* $\mathfrak{b}$ *with* $\beta(Z_0) = \frac{\pi}{3}$. *Then the orbit* $M := S(U(1)\times U(2))({\rm Exp}(Z_0))$ *of the isotropy action of* $SU(3)/S(U(1)\times U(2))$ *is a (non-totally geodesic) unstable minimal submanifold and we have* $i(M) = 30$.

Next we consider the case of $G/K = SU(6)/Sp(3)$, $H = SO(6)$ and $M = SO(6)({\rm Exp}\,Z_0)$ ($\beta_1(Z_0) = \frac{\pi}{4}$, $\beta_2(Z_0) = 0$) (the case of $n = 3$ in Example 5). Since the Satake diagram of G/K is as in Fig. 13, we have $\triangle' = \triangle$ and the multiplicity of each root of $\triangle'$ is equal to 4.



Fig. 13.

Furthermore, we have $m^V_\beta = m^H_\beta = 2$ for all $\beta\in\triangle'$ (see [Koi2]). Also, since $B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}} =$

$3B_{\mathfrak{h}}$ (see Page 187, 188 of [Hel]), we have $(\psi^* g_I)_{eL} = cB_{\mathfrak{g}}|_{\mathfrak{m}\times\mathfrak{m}} = \frac{1}{2}(B_{\mathfrak{g}}|_{\mathfrak{h}\times\mathfrak{h}})|_{\mathfrak{m}\times\mathfrak{m}} = \frac{3}{2}B_{\mathfrak{h}}|_{\mathfrak{m}\times\mathfrak{m}}$, which implies $\hat{B} = \frac{3}{2}B_{\mathfrak{h}}$. Let $\mathfrak{t}$ be a maximal abelian subalgebra of $\mathfrak{h}$ and $\triangle^{\mathfrak{h}^{\mathbf{c}}}$ be the root system of $\mathfrak{h}^{\mathbf{c}}$ with respect to $\mathfrak{t}^{\mathbf{c}}$ and define $\alpha^\sharp\,(\in \mathfrak{t})$ (resp. $\hat{\alpha}^\sharp\,(\in \mathfrak{t})$) by $\alpha(\cdot) = B_{\mathfrak{h}}(\alpha^\sharp,\cdot) = \hat{B}(\hat{\alpha}^\sharp,\cdot)$. Take $\alpha \in \triangle^{\mathfrak{h}^{\mathbf{c}}}$. Since $B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp) = \frac{1}{4}$ (see Page 188 of [Hel]) and $\hat{B} = \frac{3}{2}B_{\mathfrak{h}}$, we have $\hat{B}(\hat{\alpha}^\sharp,\hat{\alpha}^\sharp) = \frac{2}{3}B_{\mathfrak{h}}(\alpha^\sharp,\alpha^\sharp) = \frac{1}{6}$. From this relation, it follows that $\hat{B} = 12\bar{B}$. Hence we have $a_\lambda = \frac{1}{12}\bar{a}_\lambda$ for each $\lambda \in D(H)$. Since $\triangle'$ is $(\mathfrak{a}_2)$-type and $\Pi = \{\beta_1,\beta_2\}$ is a simple root system of $\triangle'_+$, we have $\triangle'_+ = \{\beta_1,\,\beta_2,\beta_1+\beta_2\}$, $l_i = \sqrt{B_{\mathfrak{g}}(\beta_i^\sharp,\beta_i^\sharp)} = \sqrt{\frac{1}{2}B_{\mathfrak{g}}(\widetilde{\beta}_i^\sharp,\widetilde{\beta}_i^\sharp)} = \frac{1}{2\sqrt{3}}$ $(i=1,2)$, $\theta_{12} = \frac{2}{3}\pi$. Also Z_0 satisfies the condition (II) in Theorem A, the constant $\hat{c}$ in Theorem A is equal to 1. Hence the symmetric matrices C_1 and C_2 in (Step II) are given by

$$C_1 = \frac{1}{144}\begin{pmatrix} 5 & -1 \\ -1 & 5 \end{pmatrix}$$

and

$$C_2 = \frac{1}{24}\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix},$$

respectively. It is shown that the maximum of the value κ such that $C_1 - \kappa C_2$ is neither a positive operator nor a negative operator is equal to $\frac{2}{3}$. Thus we have $\|\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)\|_{\mathrm{op}} = \frac{2}{3}$. Since the equivalence class μ of the complexification of the isotropy representation of $G/H(= SU(6)/SO(6))$ is $(0\,2\,0)$, we have $\bar{a}_\mu = -12$ in terms of Table 1 in [MP] and the relation (2.3) in Page 10 of [MP]. Hence we have $a_\mu = -1$. Therefore we have $a_\mu - \|\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)\|_{\mathrm{op}} = -\frac{5}{3}$. Similarly we can calculate a_λ for other $\lambda \in D(H)$. By comparing the values of a_μ and $a_\mu - \|\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)\|_{\mathrm{op}}$ with those of a_λ's ($\lambda \in D(H)$ s.t. $\lambda \neq \mu$), we have $D_1(H) = \{(0\,0\,0),\ (0\,0\,1),(0\,1\,0),(0\,0\,2),(1\,0\,1),(0\,1\,1),(0\,2\,0)\}$ and $D_2(H) = \{(0\,0\,0),\ (0\,0\,1),(0\,1\,0),(0\,0\,2),(1\,0\,1),(0\,1\,1),(0\,2\,0),(0\,0\,3),\ (1\,0\,2),(0\,1\,2),(0\,2\,1),(1\,1\,1)\}$. Since $\triangle'^V_{Z_0} = \{\beta_2\}$ and $\triangle'^H_{Z_0} = \emptyset$, we have $\mathfrak{l} = \mathfrak{z}_{\mathfrak{h}}(\mathfrak{b}) + \mathfrak{f}_{\beta_2}\cap\mathfrak{h}$. Also we have $\dim\mathfrak{l} = \dim H - (\dim G/K - \dim(\mathfrak{b} + \mathfrak{p}_{\beta_2}\cap\mathfrak{q}) = 5$. From these facts, we have $\mathfrak{l} = \mathfrak{so}(2) + \mathfrak{so}(2) + \mathfrak{su}(2)$. Hence, by using Table 2 (the branching rules) in [MP] and the branching rule for $(\mathfrak{a}_1) \subset (\mathfrak{a}_2)$ (see Page 19 ~ 20 of [MP]), we have the following table:

λ	$\lambda\|_L$	m_λ
$(0\,0\,0)$	(0)	1
$(0\,0\,1)$	$2(0)\oplus(1)$	4
$(0\,1\,0)$	$2(0)\oplus 2(1)$	6
$(0\,0\,2)$	$3(0)\oplus 2(1)\oplus(2)$	10
$(1\,0\,1)$	$4(0)\oplus 4(1)\oplus(2)$	15
$(0\,1\,1)$	$4(0)\oplus 5(1)\oplus 2(2)$	20
$\mu=(0\,2\,0)$	$3(0)\oplus 4(1)\oplus 3(2)$	20
$(0\,0\,3)$	$4(0)\oplus 3(1)\oplus 2(2)\oplus(3)$	20
$(1\,0\,2)$	$6(0)\oplus 7(1)\oplus 4(2)\oplus(3)$	36
$(0\,1\,2)$	$6(0)\oplus 8(1)\oplus 5(2)\oplus 2(3)$	45
$(0\,2\,1)$	$6(0)\oplus 9(1)\oplus 7(2)\oplus 3(3)$	60
$(1\,1\,1)$	$8(0)\oplus 12(1)\oplus 8(2)\oplus 2(3)$	64

Table 3.

Also, since $[\rho_\mu|_L]=3(0)\oplus 4(1)\oplus 3(2)$, $\mathfrak{m}^\perp=\mathfrak{b}+\mathfrak{p}_{\beta_2}\cap\mathfrak{q}$ and $\dim\mathfrak{b}=\dim\mathfrak{p}_{\beta_2}=2$, we have

$$[(\rho_\mu|_L)|_{(\mathfrak{m}^\perp)^{\mathbf{c}}}]=(0)\oplus(2).$$

Hence we have

$$\sum_{\lambda\in D_1(H)} m_\lambda \dim \mathrm{Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}})=376$$

and

$$\sum_{\lambda\in D_2(H)} m_\lambda \dim \mathrm{Hom}_L(V_{\rho_\lambda},(\mathfrak{m}^\perp)^{\mathbf{c}})=3155.$$

Therefore we obtain the following result.

Proposition 4.5. *Let $\Pi=\{\beta_1,\beta_2\}$ be the simple root system of the positive root system $\triangle_+$ of $SU(6)/Sp(3)$ and Z_0 be the element of $\mathfrak{b}$ with $\beta_1(Z_0)=\frac{\pi}{4}$ and $\beta_2(Z_0)=0$. Then the orbit $M:=SO(6)(\mathrm{Exp}(Z_0))$ of the Hermann action $SO(6)\curvearrowright SU(6)/Sp(3)$ is a (non-totally geodesic) unstable minimal submanifold and we have $376\le i(M)\le 3155$.*

Next we consider the case of $G/K=SU(8)/Sp(4)$, $H=SO(8)$ and $M=SO(8)(\mathrm{Exp}\,Z_0)$ ($\beta_1(Z_0)=\frac{\pi}{4}$, $\beta_2(Z_0)=\beta_3(Z_0)=0$) or $M=SO(8)(\mathrm{Exp}\,Z_1)$ ($\beta_2(Z_1)=\frac{\pi}{4}$, $\beta_1(Z_1)=\beta_3(Z_1)=0$) (the case of $n=4$ in Example 5). For simplicity, we set $M_i:=SO(8)(\mathrm{Exp}\,Z_i)$ $(i=1,2)$. Since the Satake diagram of G/K is as in Fig. 14, we have $\triangle'=\triangle$ and the multiplicity of each root of $\triangle'$ is equal to 4.



$(\widetilde{\beta}_i|_{\mathfrak b} = \beta_i)$

Fig. 14.

Furthermore, we have $m_\beta^V = m_\beta^H = 2$ for all $\beta \in \triangle'$ (see [Koi2]). Also, since $B_{\mathfrak g}|_{\mathfrak h\times\mathfrak h} = \frac{8}{3}B_{\mathfrak h}$ (see Page 187, 188 of [Hel]), we have $(\psi^* g_I)_{eL} = cB_{\mathfrak g}|_{\mathfrak m\times\mathfrak m} = \frac{1}{2}(B_{\mathfrak g}|_{\mathfrak h\times\mathfrak h})|_{\mathfrak m\times\mathfrak m} = \frac{4}{3}B_{\mathfrak h}|_{\mathfrak m\times\mathfrak m}$, which implies $\hat{B} = \frac{4}{3}B_{\mathfrak h}$. Let $\mathfrak t$ be a maximal abelian subalgebra of $\mathfrak h$ and $\triangle^{\mathfrak h^{\mathbf c}}$ be the root system of $\mathfrak h^{\mathbf c}$ with respect to $\mathfrak t^{\mathbf c}$ and define $\alpha^\sharp\,(\in \mathfrak t)$ (resp. $\hat{\alpha}^\sharp\,(\in \mathfrak t)$) by $\alpha(\cdot) = B_{\mathfrak h}(\alpha^\sharp, \cdot) = \hat{B}(\hat{\alpha}^\sharp, \cdot)$. Take $\alpha \in \triangle^{\mathfrak h^{\mathbf c}}$. Since $B_{\mathfrak h}(\alpha^\sharp, \alpha^\sharp) = \frac{1}{6}$ (see Page 188 of [Hel]) and $\hat{B} = \frac{4}{3}B_{\mathfrak h}$, we have $\hat{B}(\hat{\alpha}^\sharp, \hat{\alpha}^\sharp) = \frac{3}{4}B_{\mathfrak h}(\alpha^\sharp, \alpha^\sharp) = \frac{1}{8}$. From this relation, it follows that $\hat{B} = 16\bar{B}$. Hence we have $a_\lambda = \frac{1}{16}\bar{a}_\lambda$ for each $\lambda \in D(H)$. Since $\triangle'$ is $(\mathfrak a_3)$-type and $\Pi = \{\beta_1, \beta_2, \beta_3\}$ is a simple root system of $\triangle'_+$, we have $\triangle'_+ = \{\beta_1, \beta_2,, \beta_3, \beta_1+\beta_2, \beta_2+\beta_3, \beta_1+\beta_2+\beta_3\}$, $l_i = \sqrt{B_{\mathfrak g}(\beta_i^\sharp, \beta_i^\sharp)} = \sqrt{\frac{1}{2}B_{\mathfrak g}(\widetilde{\beta}_i^\sharp, \widetilde{\beta}_i^\sharp)} = \frac{1}{4}$ $(i = 1, 2, 3)$, $\theta_{12} = \theta_{23} = \frac{2}{3}\pi$ and $\theta_{13} = \frac{\pi}{2}$. The symmetric matrix C_2 in (Step II) is given by

$$C_2 = \frac{1}{32}\begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}.$$

Since Z_0 and Z_1 satisfy the condition (II) in Theorem A, the constant $\hat{c}$ in Theorem A is equal to 1. In the case of $M = M_0$, the symmetric matrix C_1 in (Step II) is given by

$$C_1 = \frac{1}{128}\begin{pmatrix} 3 & -1 & 1 \\ -1 & 3 & -3 \\ 1 & -3 & 3 \end{pmatrix}.$$

Also, in the case of $M = M_1$, it is given by

$$C_1 = \frac{1}{128}\begin{pmatrix} 2 & -1 & 0 \\ -1 & 3 & -1 \\ 0 & -1 & 2 \end{pmatrix}.$$

In both cases, it is shown that the maximum of the value κ such that $C_1 - \kappa C_2$ is neither a positive operator nor a negative operator is equal to $\frac{1}{2}$, that is, $||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}} = ||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_1}\cup\triangle'^H_{Z_1})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}} = \frac{1}{2}$. Since the equivalence class μ of the complexification of the isotropy representation of $G/H(= SU(8)/SO(8))$ is $(2\,0\,0\,0)$, we have $\bar{a}_\mu = -16$ in terms of Table 1 in [MP] and the relation (2.3) in Page 10 of [MP]. Hence we have $a_\mu = -1$. Therefore we have $a_\mu - ||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_0}\cup\triangle'^H_{Z_0})}$

$m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}} = a_\mu - ||\sum_{\beta\in\triangle'_+\setminus(\triangle'^V_{Z_1}\cup\triangle'^H_{Z_1})} m_\beta\hat{c}(\beta\otimes\beta^\sharp)||_{\mathrm{op}} = -\frac{3}{2}$. Similarly we can calculate a_λ for other $\lambda\in D(H)$. By comparing the values of a_λ's ($\lambda\in D(H)$ s.t. $\lambda\neq\mu$) with -1 and $-\frac{3}{2}$, we have $D_1(H)=\{(0\,0\,0\,0),\ (1\,0\,0\,0), (0\,1\,0\,0),\ (2\,0\,0\,0),\ (1\,0\,1\,0)\}$ and $D_2(H)=\{(0\,0\,0\,0),\ (1\,0\,0\,0), (0\,1\,0\,0),\ (2\,0\,0\,0),\ (1\,0\,1\,0),\ (1\,1\,0\,0)\}$. From $\triangle'^V_{Z_0}=\{\beta_2,\beta_3,\beta_2+\beta_3\}$ and $\triangle'^H_{Z_0}=\emptyset$, we can show $\mathfrak{l}_0:=\mathfrak{h}_{Z_0}=4\mathfrak{so}(2)+2\mathfrak{su}(2)$. Also, from $\triangle'^V_{Z_1}=\{\beta_1,\beta_3\}$ and $\triangle'^H_{Z_1}=\emptyset$, we can show $\mathfrak{l}_1:=\mathfrak{h}_{Z_1}=8\mathfrak{so}(2)$. Hence, by using Table 2 (the branching rules) in [MP] (see Page 19 $\sim$ 20 of [MP]), we have the following table:

λ	$\lambda\vert_{L_0}$	m_λ
$(0\,0\,0\,0)$	$(0-0)$	1
$(1\,0\,0\,0)$	$(1-3)$	8
$(0\,1\,0\,0)$	$2(0-2)\oplus(2-4)\oplus(0-6)$	28
$\mu=(2\,0\,0\,0)$	$(0-4)\oplus(2-2)\oplus(2-6)$	35
$(1\,0\,1\,0)$	$(1-1)\oplus(1-3)\oplus(1-5)\oplus(1-7)$ $\oplus(3-3)$	56
$(1\,1\,0\,0)$	$(1-1)\oplus 3(1-3)\oplus 2(1-5)\oplus(1-7)$ $(1-9)\oplus(3-3)\oplus(3-5)\oplus(3-7)$	160

Table 4.

Here we set $L_i:=H_{Z_i}$ $(i=0,1)$. Set $\mathfrak{m}_i^\perp=\exp^{-1}_{Z_i*}(T^\perp_{\mathrm{Exp}\,Z_i}SO(8)(\mathrm{Exp}\,Z_i))$ $(i=0,1)$. Also, since $[\rho_\mu|_{L_0}]=(2-6)\oplus(2-2)\oplus(0-4)$ and $\mathfrak{m}_0^\perp=9$, we have

$$[(\rho_\mu|_{L_0})|_{(\mathfrak{m}_0^\perp)^{\mathbf{c}}}]=(2-2).$$

Also,we have $\dim\mathfrak{m}_1^\perp=7$. Hence we have

$$\sum_{\lambda\in D_1(H)} m_\lambda \dim\mathrm{Hom}_{L_0}(V_{\rho_\lambda},(\mathfrak{m}_0^\perp)^{\mathbf{c}}) = \sum_{\lambda\in D_2(H)} m_\lambda \dim\mathrm{Hom}_{L_0}(V_{\rho_\lambda},(\mathfrak{m}_0^\perp)^{\mathbf{c}}) = 35,$$

$$\sum_{\lambda\in D_1(H)} m_\lambda \dim\mathrm{Hom}_{L_1}(V_{\rho_\lambda},(\mathfrak{m}_1^\perp)^{\mathbf{c}}) = 36470$$

and

$$\sum_{\lambda\in D_2(H)} m_\lambda \dim\mathrm{Hom}_{L_1}(V_{\rho_\lambda},(\mathfrak{m}_1^\perp)^{\mathbf{c}}) = 215670.$$

Therefore we obtain the following result.

Proposition 4.6. *Let* $\Pi=\{\beta_1,\beta_2,\beta_3\}$ *be the simple root system of the positive root system* $\triangle_+$ *of* $SU(8)/Sp(4)$*,* Z_0 *be the element of* $\mathfrak{b}$ *with* $\beta_1(Z_0)=\frac{\pi}{4}$ *and*

$\beta_2(Z_0) = \beta_3(Z_0) = 0$, and Z_1 be the element of $\mathfrak{b}$ with $\beta_2(Z_1) = \frac{\pi}{4}$ and $\beta_1(Z_1) = \beta_3(Z_1) = 0$. Then the orbits $M_0 := SO(8)(\mathrm{Exp}(Z_0))$ and $M_1 := SO(8)(\mathrm{Exp}(Z_1))$ of the Hermann action $SO(8) \curvearrowright SU(8)/Sp(4)$ are (non-totally geodesic) unstable minimal submanifolds and we have $i(M_0) = 35$ and $36470 \leq i(M_1) \leq 215670$.

References

[BCO] J. Berndt, S. Console and C. Olmos, Submanifolds and Holonomy, Research Notes in Mathematics 434, CHAPMAN & HALL/CRC Press, Boca Raton, London, New York Washington, 2003.

[Co] L. Conlon, Remarks on commuting involutions, Proc. Amer. Math. Soc. **22** (1969) 255-257.

[GT] O. Goertsches and G. Thorbergsson, On the Geometry of the orbits of Hermann actions, Geom. Dedicata **129** (2007) 101-118.

[He] S. Helgason, Differential geometry, Lie groups and Symmetric Spaces, Academic Press, New York, 1978.

[HPTT] E. Heintze, R.S. Palais, C.L. Terng and G. Thorbergsson, Hyperpolar actions on symmetric spaces, Geometry, topology and physics for Raoul Bott (ed. S. T. Yau), Conf. Proc. Lecture Notes Geom. Topology **4**, Internat. Press, Cambridge, MA, 1995 pp214-245.

[HTST] D. Hirohashi, H. Tasaki, H. Song and R. Takagi, Minimal orbits of the isotropy groups of symmetric spaces of compact type, Differential Goem. Appl. **13** (2000) 167-177.

[I] O. Ikawa, The geometry of symmetric triad and orbit spaces of Hermann actions, J. Math. Soc. Japan **63** (2011) 79-139.

[IST] O. Ikawa, T. Sakai and H. Tasaki, Orbits of Hermann actions, Osaka J. Math. **38** (2001) 923-930.

[Ki] T. Kimura, Stability of certain reflective submanifolds in compact symmetric spaces, Tsukuba J. Math. **32** (2008) 361-382.

[KT] T. Kimura and M.S. Tanaka, Stability of certain minimal submanifolds in compact symmetric spaces, Differential Geom. Appl. **27** (2009) 23-33.

[Koi1] N. Koike, Actions of Hermann type and proper complex equifocal submanifolds, Osaka J. Math. **42** (2005) 599-611.

[Koi2] N. Koike, Collapse of the mean curvature flow for equifocal submanifolds, Asian J. Math. **15** (2011) 101-128.

[Kol] A. Kollross, A Classification of hyperpolar and cohomogeneity one actions, Trans. Amer. Math. Soc. **354** (2001) 571-612.

[MP] W.G. Mckay and J. Patera, Tables of dimensions, indices, and branching rules for reresentations of simple Lie algebras, Lecture Notes in Pure and Applied Mathematics vol. 69, Marcel Dekker, Inc., New York and Baesel (1981).

[O] Y. Ohnita, On stability of minimal submanifolds in compact symmetric spaces, Compositio Math. **64** (1987) 157-189.

[PT] R.S. Palais and C.L. Terng, Critical point theory and submanifold geometry, Lecture Notes in Math. **1353**, Springer, Berlin, 1988.

[Ta] M.S. Tanaka, Stability of minimal submanifolds in symmetric spaces, Tsukuba J. Math. **19** (1995) 27-56.

[TT] C.L. Terng and G. Thorbergsson, Submanifold geometry in symmetric spaces, J. Differential Geometry **42** (1995) 665-718.

Department of Mathematics, Faculty of Science
Tokyo University of Science, 1-3 Kagurazaka
Shinjuku-ku, Tokyo 162-8601 Japan
(koike@ma.kagu.tus.ac.jp)